## Clinical Opioid Withdrawal Scale (COWS)

Times:	1	2	3	4
<i>Measure</i> 0 pulse ra 1 pulse ra 2 pulse ra		record beats per sitting or lying for 120		
Score	1	2	3	4
0 no repo 1 subject 2 flushed 3 beads	ing: over past ½ ort of chills or flus ive report of chills I or observable m of sweat on brow streaming off face	hing s or flushing oistness on face or face	ted for by room t	emperature or patient activity.
Score:	1	2	3	4
0 able to 1 reports 3 frequer	sit still difficulty sitting s nt shifting or extra	ation during assess still, but is able to aneous movemen re than a few sec	do so its of legs/arms	
Score:	1	2	3	4
1 pupils p 2 pupils r	oinned or normal cossibly larger the moderately dilate	size for room ligh an normal for roo d ly the rim of the ir	m light	
Score:	1	2	3	4
attributed 0 not pre 1 mild dif 2 patient	d to opiates withd sent ffuse discomfort reports severe d	rawal is scored iffuse aching of jo	pints/ muscles	sly, only the additional component
Score:	1	2	3	4
0 not pre 1 nasal s 2 nose ru	sent tuffiness or unus unning or tearing	g Not accounted to ually moist eyes or tears streamin		toms or allergies
Score:	1	2	3	4

## Clinical Opioid Withdrawal Scale (COWS)

0 no gas 1 stomac 2 nausea	ointestinal Up trointestinal sy ch cramps a or loose stool ng or diarrhea		≨ hour		
	e episodes of d	liarrhea or vom	iting		
Score:	1	2	3	4	
0 No trer 1 tremor 2 slight t	or observation of mor can be felt, but remor observat remor or musc	t not observed ble	hands		
Score:	1	2	3	4	
0 no yaw 1 yawnin 2 yawnin	ing Observation  Ining once or twice or three or more or several times	e during assess e times during	sment		
Score:	1	2	3	4	
0 none 1 patient 2 patient	ety or Irritabili reports increas obviously irrita so irritable or a	sing irritability oble ble anxious		assessment is diffic	cult
Score:	1	2	3	4	
0 skin is 3 piloere			rs standing up o	n arms	
Score:	1	2	3	4	
Total Scores:	1	2	3	4	_
13-2 25-3	! = mild opioid !4 = moderate !6 = moderatel i = severe with	opioid withdra y severe opio	id withdrawal		