



## Non-celiac gluten sensitivity: Time for sifting the grain

Luca Elli, Leda Roncoroni, Maria Teresa Bardella

Luca Elli, Leda Roncoroni, Maria Teresa Bardella, Center for Prevention and Diagnosis of Celiac Disease, Gastroenterology and Endoscopy Unit, Department of Pathophysiology and Transplantation, Università degli Studi di Milano. Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, 20122 Milano, Italy

**Author contributions:** Elli L, Roncoroni L and Bardella MT contributed equally to this paper.

**Conflict-of-interest statement:** The authors declare no conflict of interests.

**Open-Access:** This article is an open-access article which was selected by an in-house editor and fully peer-reviewed by external reviewers. It is distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

**Correspondence to:** Luca Elli, MD, PhD, Center for Prevention and Diagnosis of Celiac Disease, Gastroenterology and Endoscopy Unit, Department of Pathophysiology and Transplantation, Università degli Studi di Milano. Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Via F. Sforza 35, 20122 Milano, Italy. [dottorlucaelli@gmail.com](mailto:dottorlucaelli@gmail.com)  
Telephone: +39-2-55033384  
Fax: +39-2-55033664

Received: January 23, 2015

Peer-review started: January 23, 2015

First decision: March 10, 2015

Revised: April 1, 2015

Accepted: May 27, 2015

Article in press: May 27, 2015

Published online: July 21, 2015

### Abstract

In the last few years, a new nomenclature has been proposed for the disease induced by the ingestion of gluten, a protein present in wheat, rice, barley and oats. Besides celiac disease and wheat allergy, the most

studied forms of gluten-related disorders characterized by an evident immune mechanism (autoimmune in celiac disease and IgE-mediated in wheat allergy), a new entity has been included, apparently not driven by an aberrant immune response: the non-celiac gluten sensitivity (NCGS). NCGS is characterized by a heterogeneous clinical picture with intestinal and extraintestinal symptoms arising after gluten ingestion and rapidly improving after its withdrawal from the diet. The pathogenesis of NCGS is largely unknown, but a mixture of factors such as the stimulation of the innate immune system, the direct cytotoxic effects of gluten, and probably the synergy with other wheat molecules, are clues for the complicated puzzle. In addition, the diagnostic procedures still remain problematic due to the absence of efficient diagnostic markers; thus, diagnosis is based upon the symptomatic response to a gluten-free diet and the recurrence of symptoms after gluten reintroduction with the possibility of an important involvement of a placebo effect. The temporary withdrawal of gluten seems a reasonable therapy, but the timing of gluten reintroduction and the correct patient management approach are have not yet been determined.

**Key words:** Celiac disease; Gluten; Gluten-related disorders; Gluten sensitivity; Non-celiac gluten sensitivity

© **The Author(s) 2015.** Published by Baishideng Publishing Group Inc. All rights reserved.

**Core tip:** Gluten-related disorders are extremely relevant in gastroenterologic practice, and their prevalence has increased in recent decades. Besides celiac disease and allergy, the newly born non-celiac gluten sensitivity is still an obscure syndrome with an unknown pathogenesis and important clinical concerns regarding diagnosis and patient management.

Elli L, Roncoroni L, Bardella MT. Non-celiac gluten sensitivity: Time for sifting the grain. *World J Gastroenterol* 2015; 21(27): 8221-8226 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v21/i27/8221.htm> DOI: <http://dx.doi.org/10.3748/wjg.v21.i27.8221>

## INTRODUCTION

Wheat is a basic staple of the human diet, especially in Western countries where wheat-containing food dominates the menu. Since the start of agriculture in Mesopotamia about 10000 years ago, wheat has sustained mankind's development and health up to the present day<sup>[1]</sup>. This apparently contrasts with the selection of a genetic background susceptible to immune reactions against gluten [*i.e.*, celiac disease (CD)], the HLA DQ2 haplotype<sup>[2]</sup>. The development of adverse reactions against wheat and, in particular, against gluten, is supposed to be concomitant with the very moment humans started its ingestion. The first description of a patient affected by CD is ascribed to Aretaeus of Cappadocia, who in the 2<sup>nd</sup> century AD reported a case of chronic diarrhea and malabsorption. However, only at the end of the 19<sup>th</sup> century, Samuel Gee scientifically described the celiac syndrome in childhood and in the 1950s, Willem Dicke identified gluten as the environmental agent of CD<sup>[3]</sup>. Since then, the scientific community's interest concerning the effects of gluten on humans and the mechanisms underpinning gluten-induced intestinal and systemic damage has increased<sup>[4-6]</sup>. Nowadays, CD is considered an autoimmune disorder affecting about 1% of the worldwide population<sup>[7,8]</sup>. A different mechanism is involved in the allergic reaction to wheat (wheat allergy): a condition affecting 4% of the population with heterogeneous intestinal, respiratory, and cutaneous symptoms<sup>[6]</sup>. In addition to the food allergy against wheat, the other major forms of wheat allergy are baker's asthma and wheat-dependent exercise-induced anaphylaxis, which can lead to important anaphylactic reactions<sup>[9]</sup>.

Recently, a third clinical syndrome has been introduced under the generic umbrella of gluten-related disorder, or non-celiac gluten sensitivity (NCGS), which apparently involves neither the allergic nor autoimmune mechanisms or a specific genetic background, as in CD<sup>[4]</sup>. Thus, gluten or other wheat components are supposedly the environmental factors that induce a wide range of gluten-related diseases with different pathogenic pathways, which globally affect a large proportion of the population<sup>[4,10]</sup>.

## NCGS AND ITS POSSIBLE PATHOMECHANISMS

NCGS is a clinical syndrome characterized by both intestinal and extraintestinal symptoms, which are responsive to gluten (wheat) withdrawal from the diet. The mechanisms by which gluten induces symptoms in NCGS are largely unknown, but their difference from those present in CD and wheat allergy appears

clear. Although data is scant and conflicting, one can hypothesize that a multifactorial process is at the base of the NCGS symptomatic onset. The involvement of the innate immune system, a direct cytotoxic effect of gluten and gliadin, and the involvement of different target organs and other proteic components of wheat can be all implicated.

Firstly, the study by Sapone *et al.*<sup>[11]</sup> demonstrated in a series of NCGS subjects reporting different gastrointestinal symptoms that intestinal permeability and adaptive immune responses are not involved in the process. In fact, by using the lactulose/mannitol test and investigating the main components of tight junctions (zonulin and claudins), the authors found no alterations, this fact excluding the presence of a "leaky gut" as in CD or other intestinal inflammatory conditions. The concomitant duodenal increase of toll-like receptor-2 and the significant reduction of the T-regulatory cell marker FOXP3 compared to healthy subjects and CD patients suggested a stimulation of the innate immune system that does not involve the interferon  $\gamma$  pathway. In the duodenal histology of NCGS, a slight increase of  $\gamma$  and  $\delta$  intraepithelial lymphocytes was reported. The data collected on the absence of an adaptive immune response stimulation seems to be confirmed by another study that investigated the effect of a gluten-containing diet in patients affected by a diarrhea-type irritable bowel syndrome (IBS)<sup>[12]</sup>. In contrast with the previous studies, an alteration of gut permeability was reported in these patients, especially in subjects carrying a CD-compatible genetic background (*i.e.*, HLA DQ2). However, the definitive exclusion of the adaptive immune system is not possible. Based on the findings published by Brottveit *et al.*<sup>[13]</sup>, who explored and compared the initial mucosal immunologic events in CD and NCGS subjects before and after a gluten challenge, an increase of interferon  $\gamma$  and heat shock protein 27 levels after gluten challenge was evidenced together with an increased density of intraepithelial lymphocytes, this latter independent from the gluten challenge. Differently, other authors failed to identify any gliadin-related immunologic alteration in the duodenal mucosa of NCGS patients<sup>[14]</sup>.

Besides the possible alteration of the immune system, gluten presents some peculiar biologic properties that induce intestinal damage<sup>[15]</sup>. Different *in vitro* studies on two- and three-dimensional cell cultures (mainly enterocytes) demonstrated that the treatment with gluten and gliadin induces alterations of the main cellular homeostatic processes. Cellular morphology and motility<sup>[16,17]</sup> and cytoskeleton organization and intercellular contact through the tight junction proteins are all altered<sup>[18,19]</sup>. Treated cells present a reduced viability due to the stimulation of apoptosis and a reduction of synthesis of nucleic acids (DNA and RNA) and proteins<sup>[20]</sup>. Moreover, the oxidative balance is disrupted by a decrease of reduced glutathione and thiols<sup>[18,21]</sup>.

While the researchers' attention is focused on gluten, other molecules also contained in wheat may be involved in the NCGS symptoms. Among them, the amylase trypsin inhibitor enzymatic family is a known trigger of the toll-like receptor-4 pathway of the innate immune system, stimulating the release of pro-inflammatory cytokines in cells from CD and non-CD patients and biopsies<sup>[22]</sup>. Other authors have indicated that the fermentable oligosaccharides, disaccharides, monosaccharides and polyols-present in wheat (fructans) and other foods and known as FODMAPs-are important factors in causing unspecific gastrointestinal symptoms (especially bloating). These poorly absorbed short-chain carbohydrates are fermented in the colon with gas release, the resulting intestinal distension being at the base of reported symptoms. In a placebo-controlled, crossover rechallenge study, Biesiekierski *et al.*<sup>[23]</sup> found no evidence of a specific dose-dependent effect of gluten in patients with suspected NCGS undergoing a low FODMAPs diet. Although gravely by an important nocebo effect, their data suggest that NCGS might be confused with FODMAP sensitivity, and gluten might be only a cofactor (or a confounder) when looking for the origin of gastrointestinal symptoms. In a recent study, a low FODMAP diet was effective in controlling gastrointestinal symptoms in patients with intestinal bowel disease<sup>[24]</sup>, thus supporting its use as a first-line therapy. However, the data from the reported studies are extremely heterogeneous and even scant in the case of amylase trypsin inhibitor, thus making their interpretation difficult. The main bias is caused by the absence of a clear diagnostic flowchart for NCGS as all the studies simply enrolled patients clinically responding to a period of gluten-free dieting without excluding the placebo effect.

## HOW MANY PATIENTS WITH NCGS ARE THERE AND WHICH SYMPTOMS DO THEY REPORT?

Subjects with NCGS claim a variety of intestinal and/or extra-intestinal symptoms, similar to those occurring in patients affected by CD or IBS<sup>[25]</sup>. Symptoms occur after gluten ingestion; they improve/disappear when gluten is withdrawn from the diet or recur within a few hours or days on gluten challenge.

In a recent Italian, multicenter prospective survey of 486 subjects with symptomatic benefit from a gluten-free diet (62% of these subjects underwent intestinal biopsy to exclude CD), the large majority of patients reported more than two associated gastrointestinal or extraintestinal symptoms<sup>[10]</sup>. The most frequent intestinal symptoms (> 80%) were bloating and abdominal pain, while more than 50% of patients reported diarrhea, 27% alternating bowel habits, and 24% constipation. Also, epigastric pain was frequent and followed, with decreasing prevalence, by nausea, aerophagia, gastroesophageal reflux disease, and aphthous stomatitis. In this study, as in

many other studies on NCGS, systemic extraintestinal symptoms were frequently present: tiredness, lack of well-being, and neuropsychiatric symptoms such as headache, anxiety, "foggy mind", arm/leg numbness, and depression. Additional extraintestinal manifestations were as reported: muscle or joint pain, weight loss, dermatitis, and skin rash. Looking at the data regarding gluten withdrawal, NCGS is suspected in children (with fatigue apparently as their most frequent systemic symptom) and in adults, with a higher prevalence in females than in males<sup>[6,26]</sup>. Additionally, there are reports that NCGS is more common in families with a history of CD<sup>[26-29]</sup>.

There is some overlap between IBS and NCGS. In fact, in patients fulfilling the Rome III criteria for IBS, mainly in those with diarrhea, both predominant and in mixed form, NCGS has been diagnosed in a high percentage of cases: these have been sub-grouped as having gluten-sensitive IBS<sup>[30-32]</sup>. Similarly, in patients affected by allergic disorders, a high prevalence of NCGS has been reported (77/262 patients), thus confirming the need for a clearer definition of both the syndrome and the effects of gluten on the gastrointestinal tract<sup>[33]</sup>.

An objective approach to NCGS diagnosis is not available to date, as significant serologic and histologic markers are lacking; thus, the epidemiology of the NCGS disorder is quite approximate and not based on powerful studies. The reported prevalence of NCGS in the current literature varies between 0.6% and 6%. The former prevalence rate results from the National Health and Nutrition Examination Survey program, based on interviews and medical examinations carried out by general practitioners<sup>[34]</sup>, while the latter rate is the outcome of the experience by the Tertiary Center for Celiac Research at the University of Maryland<sup>[6]</sup>. At present, we should consider as affected by NCGS only those subjects who have been extensively studied to exclude CD or wheat allergy and have responded to a blind placebo-controlled gluten challenge, therefore excluding self-diagnosed self-treated subjects.

## HOW CAN WE DIAGNOSE AND TREAT NCGS?

The main problem concerning NCGS is the absence of a set of standardized diagnostic criteria. The clinical picture of NCGS seems very unspecific. Although some allergic manifestations (urticaria, asthma, labial edema, *etc.*) strongly suggest the presence of an IgE-mediated disorder, the presence of abdominal discomfort, evacuation disturbances, foggy mind, chronic fatigue, and unspecific skin manifestations do not help clinicians in any differential diagnosis. As a result, when NCGS is suspected, the first step is represented by the exclusion of other disorders, especially CD and wheat allergy. Serologic negativity to anti-transglutaminase IgA and anti-wheat component IgE antibodies and/or the absence of a duodenal atrophy at duodenal histology and skin reaction during

**Table 1** Non-celiac gluten sensitivity identikit

Name and definition	Pathogenesis	Prevalence	Diagnosis	Treatment
Non-celiac gluten/wheat sensitivity is an intestinal/extraintestinal syndrome responsive to gluten withdrawal in absence of CD and WA	Multi-factorial, including innate immunity and direct cytotoxic effects of gluten  Other molecules are potentially involved: Amylase trypsin inhibitor, FODMAPs	Ranging from 3% to 5%, though specific studies are lacking	Via placebo-controlled blind gluten challenge  Prior exclusion of celiac disease and allergy  AGA IgG, IELs, basophil activation and eosinophil duodenal/colon infiltration are potential biomarkers	No data are available for guidance  An on-demand gluten-free regimen should be considered

AGA: Anti-gliadin antibodies; CD: Celiac disease; FODMAP: Fermentable oligosaccharides, disaccharides, monosaccharides, and polyols; IELs: Intraepithelial lymphocytes; WA: Wheat allergy.

a prick test, are necessary pre-requisites.

Despite the absence of reliable NCGS biomarkers, some serologic and/or histologic alterations have been reported: (1) anti-gliadin antibodies IgG class could be present in 50% of patients with suspected NCGS<sup>[27]</sup>; (2) cytometric basophil activation is positive in 66% of patients responding to a wheat blind challenge associated to duodenal intraepithelial lymphocytosis and eosinophil infiltration of duodenum and colon<sup>[35,36]</sup>; and (3) mild duodenal intraepithelial lymphocytosis could be present in about 50% of NCGS cases, suggesting a microscopic enteritis<sup>[10,36,37]</sup>.

Once alternative causes of a patient's clinical status have been excluded, the second step is to focus on a link between symptoms and gluten ingestion with an on/off mechanism. Typically, symptoms would arise rapidly or at least after a few hours from the ingestion of gluten-containing food and would disappear "instantly" once gluten is withdrawn from the diet. This passage meets some criticism. Firstly, the period of gluten exclusion to ascertain the symptomatic response is not standardized, therefore, it is not clear how long the gluten-free diet should be maintained in absence of any clinical response. Secondly, the gluten-free dietary regimen during the proof period (if you pardon the pun) should be kept absolutely strict-the same way as for CD-to facilitate nutritional control. Thirdly, how should a gluten-free diet response be checked and measured, and how great should the symptomatic improvement be in order to define a patient responsive to gluten exclusion? Because of the absence of reliable biomarkers altered by gluten ingestion, a symptomatic trend can be measured using scores (as the gastrointestinal symptom rating scale) validated for functional diseases or visual analogue scales<sup>[38]</sup>. The quantification of symptomatic improvement is at the base of a common definition of "gluten-free diet response". Although necessary, a response to a gluten-free diet would be insufficient to define a patient as gluten sensitive, due to the possibility of a placebo effect, which is known to be

present in functional diseases<sup>[39]</sup>. Thus, a third step is required in order to eliminate or reduce the impact of the placebo effect in the diagnostic process. To reach this goal, a double- or at least single-blind challenge should be proposed to patients while they are on their gluten-free diet, to make evident the symptomatic onset during the blind assumption of gluten in a sort of n-of-1 process<sup>[40]</sup>. The problems to be solved here are: What dosage of gluten? Which vehicle should be used? How long should the challenge last? Which method to use for symptoms quantification?

## POTENTIAL MANAGEMENT OPTIONS AND CONCLUSIONS

Once the diagnosis of NCGS is reasonably reached, the management and follow-up of patients is completely obscure. A logical approach is to undertake a gluten-free dietary regimen for a limited period (*e.g.*, six months), followed by the gradual reintroduction of gluten. During the gluten-free diet, the ingestion of prolamine peptide (gliadin)-derived from wheat, rye, barley, oats, bulgur, and hybrids of these cereal grains-should be avoided. Rice, corn, and potatoes have been the typical substitutes, but nowadays other different cereals and pseudocereals, such as amaranth, buckwheat, manioc, fonio, teff, millet, quinoa, and sorghum, can be used. After some period on a gluten-free diet, the reintroduction of gluten can start with cereals of low gluten content (*e.g.*, oats). In addition, einkorn farro (*Triticum monococcum*) can be used, having no direct *in vitro* or *ex vivo* toxicity and low (7%) gluten content<sup>[41]</sup>.

Chronically, gluten-free-diet periods can be administered during symptomatic relapses as a sort of "on-demand" therapy. NCGS is a clinical syndrome hampered by many obscure sides (summarized in Table 1). Its pathogenesis and clinical presentation are not clear, and the diagnostic flowchart and patient management are not standardized. However, in the



epidemiologic dimension, the possibility to spare a large number of patients undergoing a drug-based therapy with side effects and costs is intriguing and makes NCGS a hot topic in gastroenterology.

## REFERENCES

- Rostami K, Malekzadeh R, Shahbazkhani B, Akbari MR, Catassi C. Coeliac disease in Middle Eastern countries: a challenge for the evolutionary history of this complex disorder? *Dig Liver Dis* 2004; **36**: 694-697 [PMID: 15506671]
- Lionetti E, Catassi C. Co-localization of gluten consumption and HLA-DQ2 and -DQ8 genotypes, a clue to the history of celiac disease. *Dig Liver Dis* 2014; **46**: 1057-1063 [PMID: 25200477 DOI: 10.1016/j.dld.2014.08.002]
- Losowsky MS. A history of coeliac disease. *Dig Dis* 2008; **26**: 112-120 [PMID: 18431060 DOI: 10.1159/000116768]
- Catassi C, Bai JC, Bonaz B, Bouma G, Calabrò A, Carroccio A, Castillejo G, Ciacci C, Cristofori F, Dolinsek J, Francavilla R, Elli L, Green P, Holtmeier W, Koehler P, Koletzko S, Meinhold C, Sanders D, Schumann M, Schuppan D, Ullrich R, Vécsei A, Volta U, Zevallos V, Sapone A, Fasano A. Non-Celiac Gluten sensitivity: the new frontier of gluten related disorders. *Nutrients* 2013; **5**: 3839-3853 [PMID: 24077239 DOI: 10.3390/nu5103839]
- Catassi C, Kryszak D, Bhatti B, Sturgeon C, Helzlsouer K, Clipp SL, Gelfond D, Puppa E, Sferruzza A, Fasano A. Natural history of celiac disease autoimmunity in a USA cohort followed since 1974. *Ann Med* 2010; **42**: 530-538 [PMID: 20868314 DOI: 10.3109/07853890.2010.514285]
- Sapone A, Bai JC, Ciacci C, Dolinsek J, Green PH, Hadjivassiliou M, Kaukinen K, Rostami K, Sanders DS, Schumann M, Ullrich R, Villalta D, Volta U, Catassi C, Fasano A. Spectrum of gluten-related disorders: consensus on new nomenclature and classification. *BMC Med* 2012; **10**: 13 [PMID: 22313950 DOI: 10.1186/1741-7015-10-13]
- Buscarini E, Conte D, Cannizzaro R, Bazzoli F, De Boni M, Delle Fave G, Farinati F, Ravelli P, Testoni PA, Lisiero M, Spolaore P. White paper of Italian Gastroenterology: delivery of services for digestive diseases in Italy: weaknesses and strengths. *Dig Liver Dis* 2014; **46**: 579-589 [PMID: 24913902]
- Gujral N, Freeman HJ, Thomson AB. Celiac disease: prevalence, diagnosis, pathogenesis and treatment. *World J Gastroenterol* 2012; **18**: 6036-6059 [PMID: 23155333 DOI: 10.3748/wjg.v18.i42.6036]
- Soares-Weiser K, Takwoingi Y, Panesar SS, Muraro A, Werfel T, Hoffmann-Sommergruber K, Roberts G, Halken S, Poulsen L, van Ree R, Vlieg-Boerstra BJ, Sheikh A. The diagnosis of food allergy: a systematic review and meta-analysis. *Allergy* 2014; **69**: 76-86 [PMID: 24329961 DOI: 10.1111/all.12333]
- Volta U, Bardella MT, Calabrò A, Troncone R, Corazza GR. An Italian prospective multicenter survey on patients suspected of having non-celiac gluten sensitivity. *BMC Med* 2014; **12**: 85 [PMID: 24885375 DOI: 10.1186/1741-7015-12-85]
- Sapone A, Lammers KM, Casolaro V, Cammarota M, Giuliano MT, De Rosa M, Stefanile R, Mazzarella G, Tolone C, Russo MI, Esposito P, Ferraraccio F, Carteni M, Riegler G, de Magistris L, Fasano A. Divergence of gut permeability and mucosal immune gene expression in two gluten-associated conditions: celiac disease and gluten sensitivity. *BMC Med* 2011; **9**: 23 [PMID: 21392369 DOI: 10.1186/1741-7015-9-23]
- Vazquez-Roque MI, Camilleri M, Smyrk T, Murray JA, Marietta E, O'Neill J, Carlson P, Lamsam J, Janzow D, Eckert D, Burton D, Zinsmeister AR. A controlled trial of gluten-free diet in patients with irritable bowel syndrome-diarrhea: effects on bowel frequency and intestinal function. *Gastroenterology* 2013; **144**: 903-911.e3 [PMID: 23357715]
- Brottveit M, Beitnes AC, Tollefsen S, Bratlie JE, Jahnsen FL, Johansen FE, Sollid LM, Lundin KE. Mucosal cytokine response after short-term gluten challenge in celiac disease and non-celiac gluten sensitivity. *Am J Gastroenterol* 2013; **108**: 842-850 [PMID: 23588237 DOI: 10.1038/ajg.2013.91]
- Bucci C, Zingone F, Russo I, Morra I, Tortora R, Pogna N, Scalia G, Iovino P, Ciacci C. Gliadin does not induce mucosal inflammation or basophil activation in patients with nonceliac gluten sensitivity. *Clin Gastroenterol Hepatol* 2013; **11**: 1294-1299.e1 [PMID: 23639603 DOI: 10.1016/j.cgh.2013.04.022]
- Elli L, Dolfini E, Bardella MT. Gliadin cytotoxicity and in vitro cell cultures. *Toxicol Lett* 2003; **146**: 1-8 [PMID: 14615062 DOI: 10.1016/j.toxlet.2003.09.004]
- Roncoroni L, Elli L, Bardella MT, Perrucci G, Ciulla M, Lombardo V, Tomba C, Conte D, Doneda L. Extracellular matrix proteins and displacement of cultured fibroblasts from duodenal biopsies in celiac patients and controls. *J Transl Med* 2013; **11**: 91 [PMID: 23566365 DOI: 10.1186/1479-5876-11-91]
- Roncoroni L, Elli L, Doneda L, Piodi L, Ciulla MM, Paliotti R, Bardella MT. Isolation and culture of fibroblasts from endoscopic duodenal biopsies of celiac patients. *J Transl Med* 2009; **7**: 40 [PMID: 19497109 DOI: 10.1186/1479-5876-7-40]
- Dolfini E, Elli L, Roncoroni L, Costa B, Colleoni MP, Lorusso V, Ramponi S, Braidotti P, Ferrero S, Falini ML, Bardella MT. Damaging effects of gliadin on three-dimensional cell culture model. *World J Gastroenterol* 2005; **11**: 5973-5977 [PMID: 16273608]
- Dolfini E, Roncoroni L, Elli L, Fumagalli C, Colombo R, Ramponi S, Forlani F, Bardella MT. Cytoskeleton reorganization and ultrastructural damage induced by gliadin in a three-dimensional in vitro model. *World J Gastroenterol* 2005; **11**: 7597-7601 [PMID: 16437684]
- Giovannini C, Mancini E, De Vincenzi M. Inhibition of the cellular metabolism of Caco-2 cells by prolamins peptides from cereals toxic for coeliacs. *Toxicol In Vitro* 1996; **10**: 533-538 [PMID: 20650234 DOI: 10.1016/S0887-2333(96)00042-2]
- Caputo I, Secondo A, Lepretti M, Paoletta G, Auricchio S, Barone MV, Esposito C. Gliadin peptides induce tissue transglutaminase activation and ER-stress through Ca<sup>2+</sup> mobilization in Caco-2 cells. *PLoS One* 2012; **7**: e45209 [PMID: 23049776 DOI: 10.1371/journal.pone.0045209]
- Junker Y, Zeissig S, Kim SJ, Barisani D, Wieser H, Leffler DA, Zevallos V, Libermann TA, Dillon S, Freitag TL, Kelly CP, Schuppan D. Wheat amylase trypsin inhibitors drive intestinal inflammation via activation of toll-like receptor 4. *J Exp Med* 2012; **209**: 2395-2408 [PMID: 23209313 DOI: 10.1084/jem.20102660]
- Biesiekierski JR, Peters SL, Newnham ED, Rosella O, Muir JG, Gibson PR. No effects of gluten in patients with self-reported non-celiac gluten sensitivity after dietary reduction of fermentable, poorly absorbed, short-chain carbohydrates. *Gastroenterology* 2013; **145**: 320-328.e1-320-328.e3 [PMID: 23648697 DOI: 10.1053/j.gastro.2013.04.051]
- Halmos EP, Power VA, Shepherd SJ, Gibson PR, Muir JG. A diet low in FODMAPs reduces symptoms of irritable bowel syndrome. *Gastroenterology* 2014; **146**: 67-75.e5 [PMID: 24076059 DOI: 10.1053/j.gastro.2013.09.046]
- Eswaran S, Goel A, Chey WD. What role does wheat play in the symptoms of irritable bowel syndrome? *Gastroenterol Hepatol (N Y)* 2013; **9**: 85-91 [PMID: 23983652]
- Tanpowpong P, Broder-Fingert S, Katz AJ, Camargo CA. Age-related patterns in clinical presentations and gluten-related issues among children and adolescents with celiac disease. *Clin Transl Gastroenterol* 2012; **3**: e9 [PMID: 23238134 DOI: 10.1038/ctg.2012.4]
- Volta U, Tovoli F, Cicola R, Parisi C, Fabbri A, Piscaglia M, Fiorini E, Caio G. Serological tests in gluten sensitivity (nonceliac gluten intolerance). *J Clin Gastroenterol* 2012; **46**: 680-685 [PMID: 22138844 DOI: 10.1097/MCG.0b013e3182372541]
- Volta U, De Giorgio R. New understanding of gluten sensitivity. *Nat Rev Gastroenterol Hepatol* 2012; **9**: 295-299 [PMID: 22371218 DOI: 10.1038/nrgastro.2012.15]
- Biesiekierski JR, Newnham ED, Irving PM, Barrett JS, Haines M, Doecke JD, Shepherd SJ, Muir JG, Gibson PR. Gluten causes gastrointestinal symptoms in subjects without celiac disease: a double-blind randomized placebo-controlled trial. *Am J Gastroenterol* 2011; **106**: 508-514; quiz 515 [PMID: 21224837]

- DOI: 10.1038/ajg.2010.487]
- 30 **Verdu EF**, Armstrong D, Murray JA. Between celiac disease and irritable bowel syndrome: the “no man’s land” of gluten sensitivity. *Am J Gastroenterol* 2009; **104**: 1587-1594 [PMID: 19455131 DOI: 10.1038/ajg.2009.188]
  - 31 **Verdu EF**. Editorial: Can gluten contribute to irritable bowel syndrome? *Am J Gastroenterol* 2011; **106**: 516-518 [PMID: 21378766 DOI: 10.1038/ajg.2010.490]
  - 32 **Aziz I**, Lewis NR, Hadjivassiliou M, Winfield SN, Rugg N, Kelsall A, Newrick L, Sanders DS. A UK study assessing the population prevalence of self-reported gluten sensitivity and referral characteristics to secondary care. *Eur J Gastroenterol Hepatol* 2014; **26**: 33-39 [PMID: 24216570 DOI: 10.1097/01.meg.0000435546.87251.f7]
  - 33 **Massari S**, Liso M, De Santis L, Mazzei F, Carlone A, Mauro S, Musca F, Bozzetti MP, Minelli M. Occurrence of nonceliac gluten sensitivity in patients with allergic disease. *Int Arch Allergy Immunol* 2011; **155**: 389-394 [PMID: 21346369 DOI: 10.1159/000321196]
  - 34 **DiGiacomo DV**, Tennyson CA, Green PH, Demmer RT. Prevalence of gluten-free diet adherence among individuals without celiac disease in the USA: results from the Continuous National Health and Nutrition Examination Survey 2009-2010. *Scand J Gastroenterol* 2013; **48**: 921-925 [PMID: 23834276 DOI: 10.3109/00365521.2013.809598]
  - 35 **Carroccio A**, Brusca I, Mansueto P, D’alcamo A, Barrale M, Soresi M, Seidita A, La Chiusa SM, Iacono G, Sprini D. A comparison between two different in vitro basophil activation tests for gluten- and cow’s milk protein sensitivity in irritable bowel syndrome (IBS)-like patients. *Clin Chem Lab Med* 2013; **51**: 1257-1263 [PMID: 23183757 DOI: 10.1515/cclm-2012-0609]
  - 36 **Carroccio A**, Mansueto P, Iacono G, Soresi M, D’Alcamo A, Cavataio F, Brusca I, Florena AM, Ambrosiano G, Seidita A, Pirrone G, Rini GB. Non-celiac wheat sensitivity diagnosed by double-blind placebo-controlled challenge: exploring a new clinical entity. *Am J Gastroenterol* 2012; **107**: 1898-1906; quiz 1907 [PMID: 22825366 DOI: 10.1038/ajg.2012.236]
  - 37 **Rostami K**, Aldulaimi D, Holmes G, Johnson MW, Robert M, Srivastava A, Fléjou JF, Sanders DS, Volta U, Derakhshan MH, Going JJ, Becheanu G, Catassi C, Danciu M, Materacki L, Ghafarzadegan K, Ishaq S, Rostami-Nejad M, Peña AS, Bassotti G, Marsh MN, Villanacci V. Microscopic enteritis: Bucharest consensus. *World J Gastroenterol* 2015; **21**: 2593-2604 [PMID: 25759526 DOI: 10.3748/wjg.v21.i9.2593]
  - 38 **Svedlund J**, Sjödin I, Dotevall G. GSRS--a clinical rating scale for gastrointestinal symptoms in patients with irritable bowel syndrome and peptic ulcer disease. *Dig Dis Sci* 1988; **33**: 129-134 [PMID: 3123181 DOI: 10.1007/BF01535722]
  - 39 **Shah E**, Pimentel M. Placebo effect in clinical trial design for irritable bowel syndrome. *J Neurogastroenterol Motil* 2014; **20**: 163-170 [PMID: 24840369 DOI: 10.5056/jnm.2014.20.2.163]
  - 40 **Lillie EO**, Patay B, Diamant J, Issell B, Topol EJ, Schork NJ. The n-of-1 clinical trial: the ultimate strategy for individualizing medicine? *Per Med* 2011; **8**: 161-173 [PMID: 21695041 DOI: 10.2217/pme.11.7]
  - 41 **Pizzuti D**, Buda A, D’Odorico A, D’Inca R, Chiarelli S, Curioni A, Martines D. Lack of intestinal mucosal toxicity of Triticum monococcum in celiac disease patients. *Scand J Gastroenterol* 2006; **41**: 1305-1311 [PMID: 17060124 DOI: 10.1080/003655206.00699983]

**P- Reviewer:** Bassotti G, Ciccocioppo R, Guandalini S  
**S- Editor:** Ma YJ **L- Editor:** AmEditor **E- Editor:** Liu XM





Published by **Baishideng Publishing Group Inc**

8226 Regency Drive, Pleasanton, CA 94588, USA

Telephone: +1-925-223-8242

Fax: +1-925-223-8243

E-mail: [bpgoffice@wjgnet.com](mailto:bpgoffice@wjgnet.com)

Help Desk: <http://www.wjgnet.com/esps/helpdesk.aspx>

<http://www.wjgnet.com>



ISSN 1007-9327

