

## Vitamin D for Chronic Pain

According to a comprehensive review of the clinical research evidence, helping certain patients overcome chronic musculoskeletal pain and fatigue syndromes may be as simple, well tolerated, and inexpensive as a daily supplement of vitamin D.

By Stewart B. Leavitt, MA, PhD

*This article is adapted from the author's peer-reviewed research report, Vitamin D— A Neglected 'Analgesic' for Chronic Musculoskeletal Pain, from Pain Treatment Topics. The full report, along with a special brochure for patients, is available at [www.pain-](http://www.pain-)*


Volume 8, Issue #6

Standing apart from the various other essential nutrients, vitamin D was spotlighted recently as having special therapeutic potential. This has important implications for the management of chronic musculoskeletal pain and fatigue syndromes.

During this past June 2008, news-media headlines heralded recent clinical research that revealed benefits of vitamin D for preventing type 1 diabetes,<sup>1</sup> promoting survival from certain cancers,<sup>2</sup> and decreasing the risks of coronary heart disease.<sup>3</sup> Overlooked, however, was the traditional role of vitamin D in promoting musculoskeletal health and the considerable evidence demonstrating advantages of vitamin D therapy in helping to alleviate chronic muscle, bone and joint aches, and pains of various types.

Chronic pain—persisting more than 3 months—is a common problem leading patients to seek medical care.<sup>4-7</sup> In many cases, the causes are nonspecific, without evidence of injury, disease, or neurological or anatomical defect.<sup>8,9</sup> However, according to extensive clinical research examining adult patients of all ages, inadequate concentrations of vitamin D have been linked to nonspecific muscle, bone, or joint pain, muscle weakness or fatigue, fibromyalgia syndrome, rheumatic disorders, osteoarthritis, hyperesthesia, migraine headaches, and other chronic somatic complaints. It also has been implicated in the mood disturbances of chronic fatigue syndrome and seasonal affective disorder.

Although further research would be helpful, current best evidence demonstrates that supplemental vitamin D can help many patients who have been unresponsive to other therapies for pain. Vitamin D therapy is easy for patients to self-administer, well-tolerated, and very economical.



[1]Figure 1. Vitamin

### D Metabolism

It must be emphasized that vitamin D is not a pharmaceutical analgesic in the sense of fostering relatively immediate pain relief, and expectations along those lines would be unrealistic. Because Vitamin D supplementation addresses underlying processes, it may take months to facilitate pain relief, which can range from partial to complete. Furthermore, vitamin D supplementation is not proposed as a panacea or as a replacement for other pain treatment modalities that may benefit patient care.

### Vitamin D and 'D-ficiency'

Pharmacology. Vitamin D comprises a group of fat-soluble micronutrients with two major forms: D<sub>2</sub> (ergocalciferol) and D<sub>3</sub> (cholecalciferol)<sup>10,11</sup> (see Figure 1). Vitamin D<sub>3</sub> is synthesized in the skin via exposure of endogenous 7-dehydrocholesterol to direct ultraviolet B (UVB) radiation in sunlight and is also obtained to a small extent in the diet (see Table 1). In many countries, some foods are fortified with vitamin D<sub>3</sub>, which is the form used in most nutritional supplements.<sup>14,20</sup> Vitamin D<sub>2</sub>, on the other hand, is found in relatively few foods or supplements.<sup>14</sup>

Following vitamin D synthesis in the skin or other intake, some of it is stored in adipose tissue, skeletal muscle, and many organs,<sup>19</sup> while a relatively small portion undergoes a two-stage process of metabolism (see Figure 1). First, D<sub>2</sub> and/or D<sub>3</sub> are metabolized via hydroxylation in the liver to form 25-hydroxyvitamin D, abbreviated as 25(OH)D (also called calcidiol).<sup>10,14,18,21,22</sup> This has minimal

biological activity and serum concentrations of 25(OH)D accumulate gradually, plateauing at steady-state levels by about 40 days<sup>19,23,24</sup> to 90 days.<sup>25,26</sup>

The 25(OH)D metabolite is converted primarily in the kidneys via further hydroxylation to 1,25-dihydroxyvitamin D, abbreviated as 1,25(OH)2D (also called calcitriol). It is the most important and biologically-active vitamin D metabolite with a short half-life of only 4 to 6 hours<sup>27</sup> but can remain active for 3 to 5 days.<sup>15,20</sup> A central role of vitamin D—via its active 1,25(OH)2D metabolite—is to facilitate the absorption of calcium from the intestine and help maintain normal concentrations of this vital agent. Equally important, 1,25(OH)2D sustains a wide range of metabolic and physiologic functions throughout the body.<sup>28</sup>

Vitamin D actually is misclassified as a vitamin; it may be more appropriately considered a prohormone and its active 1,25(OH)2D metabolite—with its own receptors found in practically every human tissue—functions as a hormone. These vitamin D receptors, or VDRs, may affect the function of up to 1000 different genes<sup>18,22,29,30,31</sup> helping to control cell growth or differentiation. The VDRs themselves can differ in their genetic makeup (poly-morphism) and activity, which may account for varying individual responses to vitamin D therapy.<sup>32,33</sup>

Table 1: Sources of Vitamin D

Source	Vitamin D2 and D3 Sources Vitamin D Content
	<b>Higher-Yield Natural Sources</b>
Exposure to sunlight, ultraviolet B radiation (0.5 minimal erythemal dose)†	3000-10,000 IU D3
Salmon fresh, wild (3.5 oz*)	600-1000 IU D3
Salmon fresh, farmed (3.5 oz)	100-250 IU D3
Salmon canned (3.5 oz)	300-600 IU D3
Herring, pickled (3.5 oz)	680 IU D3
Catfish, poached (3.5 oz)	500 IU D3 >
Sardines, canned (3.5 oz)	200-360 IU D3
Mackerel, canned (3.5 oz)	200-450 IU D3
Tuna, canned (3.6 oz)	200-360 IU D3
Cod liver oil (1 tsp / 0.17 oz)	400-1400 IU D3
Eastern oysters, steamed (3.5 oz)	642 IU D3
Shiitake mushrooms fresh (3.5 oz)	100 IU D2
Shiitake mushrooms sun-dried (3.5 oz)	1600 IU D2
Egg yolk, fresh	20-148 IU D3
	<b>Fortified Foods</b>
Fortified milk	60-100 IU/8 oz, usually D3
Fortified orange juice	60-100 IU/8 oz usually D3
Infant formulas	60-100 IU/8 oz usually D3
Fortified yogurts	100 IU/8 oz, usually D3
Fortified butter	50 IU/3.5 oz, usually D3
Fortified margarine	430 IU/3.5 oz, usually D3
Fortified cheeses	100 IU/3 oz, usually D3
Fortified breakfast cereals	60-100 IU/serving, usually D3
Food labels often express vitamin D content only as % of daily value, so it is usually exact amount is in International Units (IU).	
	<b>Supplements</b>
<b>Over The Counter/Internet</b>	
Multivitamin (including vitamin D)	400-800 IU vitamin D2 or D3‡
Vitamin D (tablets, capsules)	Various doses 400-50,000 IU (primarily D3)
<b>Prescription/Pharmaceutical</b>	
Vitamin D2 (ergocalciferol)	50,000 IU/capsule
Drisdol®, Calciferol®, others(vitamin D2) liquid supplements	8000 IU/mL

Rocaltrol®, Calcigex®, others (1,25[OH]2D)  
available outside US

0.25-0.5 mcg capsules  
1 mcg/mL solution

\* 1 oz = 28.3 grams = 29.6 mL; 1 IU = 40 mcg (microgram)

† A 0.5 minimal erythematol dose of UVB radiation would be absorbed after about 10-15 minutes of exposure of arms and legs to direct sunlight (depending on time of day, season, latitude, and skin sensitivity). Dark-skinned persons would require longer.

‡ Ergocalciferol on product label signifies D2; cholecalciferol signifies D3.

References: Calcitriol<sup>12</sup>; Ergocalciferol<sup>13</sup>; Holick<sup>14</sup>; Marcus<sup>15</sup>; ODS<sup>16</sup>; Singh<sup>17</sup>; Tanczos<sup>18</sup>; and Vieth<sup>19</sup>

The discovery of vitamin D receptors in many tissues besides intestine and bone—including heart, pancreas, breast, prostate, lymphocytes, and other tissues—implies that vitamin D supplementation might have applications for treating a number of disorders. These include autoimmune diseases, diabetes, cardiovascular disease, psoriasis, hypo-parathyroidism, renal osteodystrophy, and possibly leukemia and cancers of the breast, prostate, or colon.<sup>1-3,20,34-37</sup> Research is ongoing in these areas.

If vitamin D production or intake is diminished, reabsorption of vitamin D from tissue-storage reservoirs can sustain conversion to 25(OH)D and the 1,25(OH)2D metabolite for several months. However, an abundant supply of vitamin D during certain times, such as from summer sun exposure, does not deter its complete depletion during periods of lean intake, such as during winter months.<sup>19</sup>

Numerous individual factors may affect the status of vitamin D and its metabolites in the body. Synthesis of vitamin D3 in the skin in reaction to UVB exposure is a self-limiting reaction that achieves equilibrium within 20 to 25 minutes of exposure to strong sunlight in persons with white skin, with no net increase in D3 production after that.<sup>19</sup> Persons with darker skin require longer sun exposure, but the total yield in D3 is the same. Age is another limiting factor, and it is more difficult for older persons to acquire adequate vitamin D from UVB radiation.<sup>21,28</sup>

**Adequate 25(OH)D Concentration.** Most researchers agree that a minimum 25(OH)D serum level of about 30 ng/mL or more is necessary for favorable calcium absorption and good health.<sup>14,38-41</sup> Optimal 25(OH)D concentrations are considered to range from 30 ng/mL to 50 ng/mL (see Table 2).<sup>20,35,36,42-44</sup>

Most definitions of deficiency stress that circulating 25(OH)D concentrations of 15,39,45 Concentrations 15,22,39,46-48

Furthermore, levels of circulating 25(OH)D concentrations below an optimal range but above true deficiency status might also have a negative impact. This range—spanning 20 to 29 ng/mL of 25(OH)D—would constitute insufficiency.<sup>18,27</sup>

In cases of vitamin D toxicity—usually characterized by hypercalcemia and/or hyperphosphatemia—serum 25(OH)D levels are usually elevated to >150 ng/mL.<sup>14,20</sup> Some laboratories may indicate >100 ng/mL as possibly toxic, but this could be overly conservative and reported reference ranges are not always consistent from one laboratory to another.

Otherwise healthy persons at special risk include darker-skinned individuals, the obese, the elderly, and those living in northern or southern latitudes greater than 42 degrees.<sup>20,50,51</sup> Numerous factors in modern society may also contribute to the problem, including diets devoid of the relatively few foods naturally rich in, or fortified with, vitamin D; lifestyles of work or leisure spent predominantly indoors; and concerns about sun exposure with the attendant use of clothing or lotions that almost completely block UVB radiation.<sup>50</sup>

Significant numbers of males and females of all ages and races, and in all parts of the world have been found lacking in vitamin D.<sup>14,18,52-58</sup> For example, an investigation of healthy adults (18-84 years of

age) from a northern U.S. city in February found that 87% had insufficient 25(OH)D (23

Surprisingly, even persons living in the sunniest regions are not exempt. A recent study in southern Arizona found that the mean serum 25(OH)D concentration in 637 randomly-selected subjects was about 26 ng/mL; half of them had concentrations 49

## Associations of Vitamin D and Pain

Vitamin D in Chronic Musculoskeletal Pain. From the perspective of vitamin D involvement in musculoskeletal pain, the process is presumed to begin with a lack of circulating calcium (hypocalcemia) due to inadequate vitamin D, and this sets in motion a cascade of bio-chemical reactions negatively affecting bone metabolism and health. Even mild hypocalcemia results in an elevation of parathyroid hormone (PTH) that can diminish bone density (osteopenia) and/or more severely affect bone architecture (osteoporosis).<sup>35,36</sup>

The effect relating more closely to musculoskeletal aches and pains is that increased PTH levels also impair proper bone mineralization causing a spongy matrix to form under periosteal membranes covering the skeleton. This gelatin-like matrix can absorb fluid, expand, and cause outward pressure on periosteal tissues, which generates pain since these tissues are highly innervated with sensory pain fibers.<sup>27,44,59</sup>

This dysfunction of bone metabolism (osteomalacia) is proposed in the literature as an explanation of why many patients with vitamin D inadequacies may complain of dull, persistent, generalized musculoskeletal aches, pains, and weakness.<sup>20</sup> Therefore, experts recommend that vitamin D deficiency and its potential for associated osteomalacia should be considered in the differential diagnosis of all patients with chronic musculoskeletal pain, muscle weakness or fatigue, fibromyalgia, or chronic fatigue syndrome.<sup>27</sup>

Table 2: Defining Adequate 25(OH)D Concentrations

25(OH)D Concentrations  
Major Metabolite of Vitamin D from D2 or D3

Deficient <20 ng/mL

Insufficient 20–29 ng/mL

Optimal Range 30–50 ng/mL

Potentially Toxic >150 ng/mL

In some literature 25(OH)D is expressed as nmol/L. The conversion is  
1 nmol/L = 0.4 ng/mL.

References: Heaney<sup>39</sup>; Heaney et al<sup>42</sup>; Holick<sup>14</sup>; Register et al<sup>27</sup>; Vieth<sup>24</sup>


Table 3. Vitamin D Research in Musculoskeletal Pain & Fatigue Syndromes			
Researchers/Population	Subjects	Vitamin D Status	Comments
Turner et al <sup>17</sup> - Mainz Clinic - retrospective case-series in patients with chronic musculoskeletal pain of 65 women osteopaths, mean age 47.5 years. [Reprinted often by Heston et al <sup>18</sup> ]	n=767 71% female 76% white	70% <10 ng/mL 25(OH)D mean = 15.7 ng/mL Overall mean = 16.7 ng/mL (reflecting insufficient)	Low back pain (n=77) and fibromyalgia (n=14) were the most prevalent diagnoses. More than half (50%) of all patients were using opioids (mean morphine equivalent for nighting) for an average of 17 months. Patients with deficient vitamin D were taking opioids longer, and also reported poorer physical functioning and overall health perceptions.
Hicks et al <sup>19</sup> - Italy - prospective study in elderly persons, mean age 74.5, with pain in lower extremities or back, or both.	n=958 55% females	22% <10 ng/mL 25(OH)D 50% <20 ng/mL median = 18 ng/mL, males median = 14 ng/mL, females (25(OH)D was significantly lower in older patients)	A low threshold of <10 ng/mL 25(OH)D was defined as deficiency, although, more than half of all subjects were at <20 ng/mL. Pain was associated to a greater degree with inadequate 25(OH)D for lower back pain in women than in men. Lower 25(OH)D also was associated with more depressive symptoms, poorer cognition, and higher PTH levels.
Loft et al <sup>20</sup> - Egypt - female patients with chronic (>3 mo.) low back pain compared with a pain-free control group.	60 patients 20 controls	pts 62% <10 ng/mL 25(OH)D cts 60% <10 ng/mL (this is a much higher threshold for defining deficiency than used in other studies)	25(OH)D was significantly lower (p < .05) and PTH higher (p < .05) in patients with pain. Aspartate phosphatase higher in patients with pain, no differences in calcium or phosphorus. Low sun exposure (duration and skin exposed) accounted for 98% of the variance between groups, even in this sunny climate.
Armstrong et al <sup>21</sup> - United States - patients with fibromyalgia syndrome.	n=70	56% = 10-20 ng/mL 25(OH)D 13% <10 ng/mL	96% of total considered deficient in 25(OH)D. There also was a correlation between deficient vitamin D and greater anxiety and depression.
Kiuding <sup>22</sup> - Kansas - women with breast cancer experiencing joint pain prior to cancer treatment.	Not stated.	71% vitamin D deficient (no cholecalciferol given)	In this pilot study, supplemental vitamin D (prior and unless) reduced joint pain as well as fatigue.
de la Jara et al <sup>23</sup> - Switzerland - 16 males from various countries of origin with chronic back pain and/or multiple somatic pain symptoms (consistent with osteoarthritis diagnosis).	n=33; mean age=39	97% <10 ng/mL 25(OH)D (mean=4.5 ng/mL) 47% had hypocalcemia 32% hypophosphatemia	With vitamin D supplementation, symptoms disappeared after 2.84 months in two-thirds (22/33) of patients, another 13% (5/33) had partial resolution. Mean number of Pn analgesic drugs taken declined from 5.27 to 0.85.
Coetse and Davis <sup>24</sup> - Michigan - randomly selected patients in a pain clinic with arthritis, pelvic pain, failed back surgery, and fibromyalgia.	n=56 81% female	96% <30 ng/mL 25(OH)D 81% = deficient (undefined) 50% <17 ng/mL	Patients ranged in age from 26 to 81 years; no significant differences in 25(OH)D outcomes across age groups or between female and male.
Benson et al <sup>25</sup> - Australia - Aboriginal patients with muscle pain, compared with a pain-free control group from the same population.	8 patients 8 controls	100% pts = 20 ng/mL 25(OH)D 12% cts = 20 ng/mL	Mean 25(OH)D in patients was 16 ng/mL, vs 23.3 ng/mL in controls (p = 0.17). All subjects were urban dwelling and had left their prior outdoor lifestyle with its ample exposure to sunlight.
Hillward et al <sup>26</sup> - England - South Asian women with either widespread widespread pain or specific rheumatic diagnosis.	n=100	70% = low 25(OH)D levels (see)	In 4.7% of patients for whom combined data were available, PTH was elevated and 25(OH)D was deficient - which are biomarkers of osteoporosis. Few had abnormal Ca, PO <sub>4</sub> , or ALP.
Ertal et al <sup>27</sup> - Germany - comparison of German control group vs Turkish patients with bone/muscle pain living in Germany or Turkey.	882 patients 121 controls	pts: n mean 15.7 ng/mL 25(OH)D 75% <20 ng/mL 25(OH)D cts = mean 27.4 ng/mL	Age range was 10-89 years, 47% females. There was a strong correlation between low 25(OH)D and higher rates and longer duration of generator bone and/or muscle aches and pains (often diagnosed as fibromyalgia). Lack of sun exposure, higher latitude, and female sex were important predictors for low 25(OH)D.
Machirana et al <sup>28</sup> - Scotland - South Asian young women with widespread pain.	n=114	3.5 times greater risk of <10 ng/mL 25(OH)D in those with pain than without.	This was a subset of patients from a study of 3135 South Asian subjects who demonstrated a 60% greater rate of widespread pain than their non-Asian counterparts in the UK.

 [Click to enlarge](#)

[2]

**Table 3. Vitamin D Research in Musculoskeletal Pain & Fatigue Syndromes (continued)**

Researchers/Population	Subjects	Vitamin D Status	Comments
Staker et al <sup>71</sup> - Boston VA Medical Center - patients with painful and radiographically confirmed knee osteoarthritis.	n=221; mean age=67	48% > 20 ng/mL 25(OH)D	Subjects with lower vitamin D had more pain and disability and were weaker. In patients with increasing vitamin D levels during a 30 mo. monitoring period there were corresponding improvements in disability and pain scores.
Hague et al <sup>72</sup> - Johns Hopkins, Baltimore - general rheumatology patients with pain	n=48; 87% female; mean age=59	94% > 10-20 ng/mL 25(OH)D 12% < 10 ng/mL	A total 92% were vitamin D deficient. The most common diagnoses were rheumatoid arthritis, inflammatory polyarthritis, chronic musculoskeletal pain, and polymyalgia rheumatica.
van der Heijden et al <sup>73</sup> - Netherlands: females with progressive muscle weakness and pain >4 mo. [case report]	n=3	Low 25(OH)D (undetectable)	Also had decreased phosphate, increased alkaline phosphatase. After vitamin D supplementation, pain resolved and muscle strength improved "within a week."
Block <sup>74</sup> - Maine - white patients with chronic widespread musculoskeletal pain, 68% diagnosed as fibromyalgia.	n=104 85% female	47% > 20 ng/mL 25(OH)D 9% < 10 ng/mL	The author alleged that these levels were not sufficiently deficient to account for the pain syndromes in these patients, but this belief is not consistent with other research. PTH levels were not measured.
Potkinoff and Quigley <sup>75</sup> - Minnesota - patients with chronic, nonspecific musculoskeletal pain (excluded fibromyalgia, complex regional pain syndrome, other disorders)	n=180	92% < 13 ng/mL 25(OH)D 100% < 20 ng/mL 28% < 8 ng/mL	Age range up to 65 years, with all age groups affected. Overall, no significant differences between males and females. Darker-skinned patients had greater 25(OH)D deficiencies and deficits in all patients were more pronounced in winter.
Al Faraj and Al Mutairi <sup>76</sup> - Saudi Arabia - patients with idiopathic chronic (>6 mo.) back pain (probable osteoarthritis, as patients with diagnosed anatomical, neuropathic, or mechanical causes were excluded)	n=360 50% female	67% > 10 ng/mL 25(OH)D (Low threshold for "normal" defined as >38 ng/mL)	Age range up to 52 years. After vitamin D supplementation for 3 months symptom improvement was seen in 95% of all patients and in 100% of those who would be considered as severely 25(OH)D deficient pretreatment.
Halsonen et al <sup>77</sup> - Canada - female patients with fibromyalgia.	n=26	48% < 20 ng/mL 25(OH)D (Group mean = 26 ng/mL)	PTH elevation also was evident in those with deficient 25(OH)D.
Grang et al <sup>78</sup> - Denmark - female Arab patients with bone pain and muscle pain and weakness compared with pain-free female Danish controls.	55 patients 22 controls	46% 85% < 3 ng/mL 25(OH)D 96% < 8 ng/mL also = mean 19 ng/mL	High dose vitamin D supplementation increased 25(OH)D 5-fold in 3 months and was paralleled by significant reductions in muscle and bone discomfort. Pretreatment, patients had elevated PTH but only 8% had subnormal calcium; alkaline phosphatase was normal. Note: most symptom-free controls had 25(OH)D levels that could be considered inadequate.
Prabhala et al <sup>79</sup> - SUNY, New York - aches, pains, and severe myopathy in patients confined to wheelchairs (case report).	n=5	Confirmed low 25(OH)D (undetectable), PTH elevated.	Weakness had been attributed to old age, diabetic neuropathy, or general debility; vitamin D supplementation resolved body aches and pains, and restored normal muscle strength in 4-6 weeks - 4 patients became fully mobile and had normal 25(OH)D.
McLinton et al <sup>80</sup> - United States - osteoarthritis of the knee recorded in persons participating in the Framingham Study.	n=75-knees (87% progressively worsening)	33% < 14 ng/mL 25(OH)D 50% < 35 ng/mL	73% of cases involved vitamin D intake < 400 IU/day. Low intake and low 25(OH)D resulted in a 3-fold risk of progressive osteoarthritis; although, there was no evidence of low intake/low 25(OH)D as causing osteoarthritis in normal knees. Conclusion was that persons with osteoarthritis and 25(OH)D < 30 ng/mL should have increased vitamin D intake.
Guth et al <sup>81</sup> - Johns Hopkins, Baltimore: hypomagnesemia (nonspecific) responsiveness to physical stimuli on response to analgesics (case report)	n=5	Range 3.2-41 ng/mL 25(OH)D	Pain resolved in 5-7 days after high dose vitamin D supplementation. In 1 patient, 25(OH)D again became deficient and pain returned, but was followed with further supplementation.



[3]

Table 3 summarizes 22 clinical investigations of vitamin D in patients with chronic musculoskeletal-related pain. These studies were conducted in various countries and included approximately 3,670 patients representing diverse populations and age groups.

The percentage of patients with pain having inadequate vitamin D concentrations ranged from 48% to 100%, depending on patient selection and the definition of 25(OH)D "deficiency." In most cases, 54,66,68,76

An interesting feature is that many of the control-group patients also had 25(OH)D concentrations that would be considered at the least as insufficient. This would more genuinely reflect the background prevalence of vitamin D inadequacy in the various populations studied, and it also confirms that vitamin D deficits do not always result in musculoskeletal pain in all persons.

Researchers also reported results of vitamin D supplementation therapy in response to the symptoms putatively related to the 25(OH)D deficiencies that they detected:

- In one large study of 360 female patients with chronic back pain, vitamin D therapy produced symptomatic improvement in 96% of all patients and in 100% of those with the most severe 25(OH)D deficiencies.<sup>74</sup> This study was of interest because only cases of idiopathic pain probably

associated with osteomalacia were included; patients with pain diagnosed as due to anatomical, neuropathic, or injury-related causes were excluded.

- In a study of 33 patients with chronic back pain and/or multiple somatic pain symptoms, researchers reported that vitamin D therapy led to a resolution of all symptoms in two-thirds of the subjects.<sup>65</sup> Partial pain relief was achieved in 18% of patients and 16% were not helped.
- A case-series report noted that aches, pains, and extreme muscle weakness were resolved by vitamin D supplementation in five patients who had been confined to wheelchairs.<sup>77</sup> Four of the patients reportedly became fully mobile upon normalization of their 25(OH)D concentrations.
- An earlier case report of five patients by Gloth and colleagues<sup>79</sup> noted that vitamin D supplementation produced significant and rapid improvements of hyperesthesias (nonspecific oversensitivity to physical stimuli) that had been unresponsive to analgesic therapy.

Taken as a whole, the research evidence supports vitamin D supplementation as a potentially important therapy for helping to ameliorate chronic, nonspecific musculoskeletal pain, and fatigue syndromes. In fair balance, however, it must be noted that randomized controlled trials (RCTs) have not been conducted to assess vitamin D therapy for pain. Such investigations would be helpful and could be important for defining which disorders are relieved most by vitamin D and the dosing regimens that are optimal for particular groups of patients. Furthermore, it is evident that not all persons with inadequate 25(OH)D levels develop musculoskeletal maladies and the specific predisposing factors affecting some persons but not others are unknown.

**Beyond Musculoskeletal Pain.** Since vitamin D receptors have been identified in skeletal muscle, myopathy—presenting as decreased muscle strength, usually in lower limbs, or merely as fatigue—is also part of the D-deficiency/osteomalacic symptom complex and may appear prior to any pain.<sup>31,76,80-83</sup> Vitamin D deficiency also has correlated with increased body sway<sup>84</sup> and an increased risk of falls that often result in painful fractures.<sup>85-89</sup> Various investigations have demonstrated benefits of vitamin D supplementation for reversing myopathy and increasing physical endurance.<sup>87,90</sup>

Clinical researchers have also found that the role of vitamin D extends beyond bone and muscle involvement in chronic pain syndromes. For example, vitamin D receptors have been identified in various brain structures, the spinal cord, and sensory ganglia.<sup>31</sup> Accordingly, results of some studies suggest non-musculoskeletal benefits of vitamin D supplementation, as follows:

- **Neuropathy.** A recently reported prospective study of 51 patients with type 2 diabetes and associated chronic, painful neuropathy found that conservative vitamin D supplementation (about 2000 IU/day) for 3 months resulted in nearly a 50% decrease in pain scores.<sup>91</sup>
- **Inflammation.** Clinical research indicates that vitamin D supplementation modulates or decreases pro-inflammatory cytokines (eg, C-reactive protein, interleukin 6 and 12, and tumor necrosis factor-alpha) while increasing anti-inflammatory cytokines (eg, interleukin-10).<sup>90,92-94</sup> Investigators have further suggested that vitamin D may help to moderate painful chronic inflammatory autoimmune conditions that are influenced by excessive cytokine activity, such as inflammatory bowel disease.<sup>18</sup>
- **Migraine Headaches.** There have been case reports of vitamin D combined with calcium supplementation to alleviate migraine headaches in postmenopausal<sup>95</sup> and premenopausal<sup>96</sup> women. Reductions in both frequency and intensity of migraines were achieved within 2 months.
- **Affective (Mood) Disturbances.** In patients with fibro-myalgia syndrome, pain, depression, and anxiety were found to be strongly associated with insufficient vitamin D.<sup>63</sup> Furthermore, large studies examining older persons (aged >65 years) found significant associations between 25(OH)D deficiencies and depressive disorders.<sup>53,62</sup> In one of the studies, inadequate 25(OH)D and depression also were highly correlated with chronic lower-back pain specifically in female patients.<sup>62</sup>

There is some evidence that seasonal affective disorder (SAD) is influenced by diminished stores of vitamin D, which would be expected to occur most commonly during winter months. In a study of



supplemental vitamin D versus broad-spectrum light therapy (phototherapy), which is often recommended for patients with SAD, vitamin D produced significant improvements in all outcome measures of depression, whereas the phototherapy group showed no significant changes.<sup>97</sup>

In a randomized, placebo-controlled study that included patients with clinical depression, those administered supplemental vitamin D had significantly enhanced mood and a reduction in negative-affect symptoms.<sup>98</sup> In similar investigations, more adequate vitamin D concentrations were associated with better physical, social, and mental functioning as measured by quality-of-life assessment instruments,<sup>99</sup> and improved scores on an assessment of well-being.<sup>100</sup>

In sum, while further research is needed, the potential benefits of vitamin D supplementation may be expanded from its role in supporting bone and muscle health to that of a complex hormonal system benefiting other conditions often associated with chronic pain.<sup>31</sup>

## Assessing Vitamin D Status

**Clinical Indicators.** From a clinical perspective, a number of factors may suggest that chronic musculoskeletal pain and related problems may be due to inadequate vitamin D intake. Researchers have stressed that the “gold standard” for a presumptive diagnosis of inadequate vitamin D is a review of patient history, lifestyle, and dietary habits that might pose risks for deficiency.<sup>101</sup> Along with this, indicators of defects in bone metabolism may include chronic muscle, bone, or joint pain, as well as persistent muscle weakness, fatigue, and possibly difficulty walking.<sup>20,101,102</sup> Radiological changes potentially associated with osteomalacia are seen only in advanced stages.<sup>20,103</sup>

Signs/symptoms of calcium deficiency (hypocalcemia) due to vitamin D deficiencies relate to neuromuscular irritability. Patients sometimes complain of paresthesias (numbness, tingling, prickling, or burning) in their lips, tongue, fingertips, and/or toes, along with fatigue and anxiety. Muscles can be painfully achy, progressing to cramps or spasms.<sup>104-107</sup> Lethargy, poor appetite, and mental confusion may be part of the syndrome.<sup>104</sup>

The diverse signs and symptoms may be erroneously attributed to other causes. Holick<sup>14,44</sup> and others<sup>51,76</sup> caution that osteomalacia due to vitamin D deficiency can be misdiagnosed as chronic fatigue syndrome, arthritis or rheumatic disease, depression, or fibromyalgia.

**Biochemical Markers.** Laboratory assessments usually pertain to the measurement of biomarkers that could denote osteomalacic processes, including:<sup>108</sup>

- Serum 25(OH)D, total serum calcium (Ca), and phosphate (PO) are decreased to below normal ranges;
- Parathyroid hormone (PTH) and total alkaline phosphatase (ALP) are elevated.

It must be understood that there are limitations to the various laboratory assays in terms of their accuracy and/or helpfulness in making or confirming a diagnosis. Assessing ALP (a surrogate marker for bone turnover), PO, or Ca are not reliable predictors of inadequate 25(OH)D concentrations or underlying osteomalacic processes.<sup>28,101</sup> Up to 20% of patients with 25(OH)D deficiency and elevated PTH may have normal Ca, PO, and ALP levels.<sup>74,108</sup>

It is generally believed that elevations of PTH may be a suitable biomarker of histological osteomalacia since, at the least, secondary hyperparathyroidism seeks to correct calcium deficits via bone resorption. A diagnosis of inadequate vitamin D with osteomalacic involvement to some extent could be presumed if PTH is elevated in association with low calcium levels, which also would serve to exclude patients with primary hyperparathyroidism due to other causes.<sup>108</sup>

Various assays for determining 25(OH)D concentrations in serum have relatively recently become available from commercial laboratories.<sup>39,109</sup> Circulating 25(OH)D reflects both D2 plus D3 intake, but not 1,25(OH)2D concentrations. Measuring 1,25(OH)2D is not recommended because it can be a poor or misleading indicator of overall vitamin D status.<sup>14,28,107</sup>

There are some concerns about the validity and utility of 25(OH)D assays, which also can be relatively expensive.<sup>108</sup> Several testing methods are available and there can be significantly large differences in results from one laboratory to the next, as well as stark variations across types of assays.<sup>21,39,46,110</sup> Error rates can be high and the reference ranges reported may be confusing or unhelpful. These concerns should not completely deter testing for biomarkers related to vitamin D deficiency. Rather, informed healthcare providers need to consider test limitations and their objectives in using such measures, keeping in mind that patient-centered care focuses on individual needs rather than relying solely on laboratory values for guidance.

While severe deficiencies in 25(OH)D and unambiguous clinical signs of osteomalacia relate most clearly to musculoskeletal pain, less severe vitamin D inadequacies are often unrecognized but nevertheless contributing sources of nociception in patients with chronic pain.<sup>60</sup> This has been proposed as a “subclinical” effect of vitamin D inadequacy, since subjective musculoskeletal pain or weakness develops prior to the emergence of more objective clinical indicators.<sup>54,76,111</sup>

Lotfi et al<sup>54</sup> proposed that in some persons even slight deficits of 25(OH)D can produce secondary hyperparathyroidism to a degree that manifests as musculoskeletal pain and/or weakness. Although vitamin D inadequacy may not be extreme enough to produce clinically diagnosable osteomalacia, it can still cause enough PTH elevation to generate increased bone turnover and loss, increased risk of microfractures, and pain or myalgia.

As Holick<sup>112</sup> has suggested, either the level of 25(OH)D is adequate for the individual patient, or it is not. If it is inadequate, subclinical osteomalacia along with multiple forms of chronic pain and myopathy may emerge and the extent of the 25(OH)D deficit in nanograms-per-milliliter may not matter – as long as the shortfall is corrected to the extent necessary.

## **Vitamin D Therapy**

**Vitamin D Intake in Healthy Persons.** In 1997, the U.S. Institute of Medicine determined that there was insufficient data to specify a Recommended Daily Allowance (RDA) for vitamin D. Instead, the organization developed very conservative Adequate Intake (AI) values of 200 IU to 600 IU per day of vitamin D, based on an assumption that as people age they would need extra vitamin D supplementation<sup>16,113</sup> (see Table 4).

According to the most recent 2005 Dietary Guidelines for Americans from the U.S. government,<sup>114</sup> and expert recommendations,<sup>14,18,24,42,58</sup> healthy children and adults of any age should consume no less than 1000 IU/day of vitamin D3 to reach and maintain minimum serum 25(OH)D concentrations of at least 30 to 32 ng/mL. However, this may be inadequate.

Clinical research trials have demonstrated that 1000 IU/day of vitamin D3 produces only modest increases in 25(OH)D that can be inconsequential for achieving and maintaining optimal concentrations of 30 ng/mL or more in some persons.<sup>26,29</sup> Vieth and colleagues<sup>26</sup> demonstrated that 4000 IU/day D3 could be safe and more effective in producing desired outcomes. Several general principles emerge from the accumulated research to date<sup>19,21,23,25,26,39,46,100,115,116,117,118</sup>:

- Concentrations of 25(OH)D are not increased in direct proportion to the amount of supplementation increase. For example, tripling the D3 dose—such as going from 600 IU/day to 1800 IU/day—does not increase the concentration of 25(OH)D by threefold.

- Any increase in 25(OH)D is also dependent on the concentration of this metabolite at the start of treatment. At equivalent vitamin D doses, patients with more severe baseline inadequacies will have larger, more rapid increases in 25(OH)D concentrations.
- What might be considered large doses of vitamin D3 by some practitioners do not produce proportionately large increases in 25(OH)D concentrations, depending on the amount of dose and duration of administration. For example, a single 50,000 IU dose of D3 may produce a significantly smaller increase in 25(OH)D than 2000 IU given daily over time, and the increases from either of these could be modest.
- However, it must be noted that continuous megadoses of vitamin D (eg, >50,000 IU) could produce robust, possibly toxic, increases in 25(OH)D concentrations over time.

In everyday practice, exceptionally large daily doses of vitamin D would rarely be recommended to patients. Researchers have noted that raising 25(OH)D from 20 ng/mL to the more optimal 30+ ng/mL range in otherwise healthy patients would require ongoing daily supplementation of only about 1300 IU<sup>39</sup> to 1700 IU<sup>119</sup> of vitamin D3. However, it should be noted that the daily adequate intake of vitamin D for maintaining health is, in most cases, lower than the amount needed as therapy for patients with chronic pain.

Table 4. Recommended Vitamin D Intake in Healthy Persons<sup>1997</sup> — Institute of Medicine

200 IU/d – children and adults to age 50 years  
 400 IU/d – men and women aged 50-70 years  
 600 IU/d – those older than 70 years

2005 — Dietary Guidelines for Americans  
 1000 IU/d – children and adults

*Note: In some of the literature International Units are used. 1 IU = 0.025 mcg or 1 mcg = 40 IU.*

Putting “D” Into Clinical Practice. In patients with pain, researchers have examined daily vitamin D supplementation ranging from 600 IU to 50,000 IU,<sup>19,27,116</sup> as well as much larger amounts. Because vitamin D has a long half-life and can take several months to reach steady-state levels, one approach to supplementation has been to administer oral or intramuscular megadoses on an infrequent basis. For example, single doses of 300,000 IU D2 have been used in the expectation that they might suffice for many months.<sup>120</sup> Hathcock et al<sup>121</sup> and others<sup>122</sup> noted that amounts of vitamin D up to 100,000 IU would not be toxic if restricted to one administration every four months, or daily for a single period of four days.

In one study of patients with chronic back pain, subjects were treated for 3 months with either 5000 IU/day or 10,000 IU/day of vitamin D3 (heavier patients >50 kg received the larger dose),<sup>74</sup> There were no adverse effects reported, and pain symptoms were relieved in 95% of the patients.

Despite the reported successes of larger-dose vitamin D supplementation, many healthcare providers may be uncomfortable with recommending such doses for their patients. And, unless pathways of vitamin D metabolism are impeded (eg, due to liver or renal disease or an interacting drug), such high doses could be unnecessary, at least as initial therapy.

In patients with chronic pain, Gloth and colleagues<sup>79</sup> observed that symptom relief often can be achieved with relatively modest increases in 25(OH)D and 1,25(OH)2D concentrations. This is possibly because the vitamin D metabolites are being rapidly consumed at tissue sites and also becoming depleted in storage depots, so they cannot accumulate in needed quantities and so any added amount is beneficial. These researchers also suggested that pain syndromes may affect vitamin D receptors, causing them to become altered in function or increased in quantity (upregulated) and, thereby, physiologically requiring extra amounts of the 1,25(OH)2D hormone.

Table 5. Vitamin D Supplementation for Chronic Pain

Proposed Conservative Dosing Protocol

- In patients with chronic, nonspecific musculoskeletal pain and fatigue

be expected that vitamin D intake from combined sources is inadequate if serum 25(OH)D are insufficient or deficient.

- All patients should take a multivitamin to ensure at least minimal intake of essential nutrients, including calcium and 400 IU to 800 IU of vitamin D.
- Recommend a daily 2000 IU vitamin D3 supplement, bringing total daily intake to 2800 IU/day (incl. from multivitamin). Extra calcium may not be necessary if intake is insufficient and/or there are concerns about osteoporosis (e.g., in postmenopausal women and elderly).
- Monitor patient compliance and results for up to 3 months. Other treatments for pain and progress do not necessarily need to be discontinued.
- If results are still lacking after 3 months, or persistent 25(OH)D deficiency is verified, consider a brief course of prescribed high-dose vitamin D3 with calcium as appropriate, followed by ongoing supplementation as maintenance.

A proposed conservative dosing protocol is outlined in Table 5. This involves adding a daily supplement of 2000 IU of vitamin D3 to a daily multivitamin regimen, bringing the total daily vitamin D3 intake to 2400 IU to 2800 IU. This is a convenient supplement dose, since inexpensive 1000 IU D3 tablets or capsules are readily available and some outlets are now stocking 2000 IU/dose tablets. The daily cost of the supplement is typically US \$ 0.10 or less.

Vitamin D3 products are preferred since they cost no more than D2 and most research indicates that D3 is more effective<sup>19,115,118,123</sup> Vieth<sup>19</sup> has strongly urged that “all use of vitamin D for nutritional and clinical purposes should specify cholecalciferol, vitamin D3.”

Vitamin D therapy would be contraindicated in patients with pre-existing excessive levels of calcium (hypercalcemia or hypercalcuria), and special caution might be advised in those prone to forming kidney stones or other calcifications.<sup>12,13,121</sup> Besides renal or hepatic dysfunction, intestinal malabsorption due to age, irritable bowel syndrome, Crohn’s disease, or celiac disease also may limit response to vitamin D therapy.<sup>20,51,124</sup>

Current therapies for chronic pain, started prior to initiating vitamin D3 therapy, do not need to be discontinued; however, it must be accepted that it could be difficult to attribute improvements to one therapy over another. This would be confounded further if new therapies for pain are started during vitamin D3 supplementation and before enough time has elapsed to evaluate its effectiveness.

If there are no improvements after several months of the proposed conservative vitamin D3 dosing protocol, more time rather than increased doses may be necessary for vitamin D3 therapy to effectively raise 25(OH)D concentrations, lower PTH levels, and/or saturate vitamin D receptors with the 1,25(OH)2D metabolite. In some cases, the patient might benefit from a course of high-dose vitamin D3 supplementation followed by more conservative doses for ongoing maintenance.

There is always the possibility that a particular chronic pain condition cannot be alleviated by vitamin D3 supplementation alone. For example, there may be a previously undetected anatomic defect, disease, or other pathology that would benefit from another type of therapeutic intervention in addition to vitamin D therapy.

At the doses recommended in Table 5, excessive accumulation of 25(OH)D or toxicity over time would not be expected and this supplementation might be continued indefinitely. It must be noted, however, that clinical investigations have largely observed subjects during months rather than years of ongoing supplementation, and long-term effects of vitamin D therapy on functions at the cellular level are still under investigation.<sup>30</sup> If there are concerns, after a year or longer the supplement might be continued at a reduced dose; it can always be increased again if pain symptoms return.

**How Long Until Improvement?** In anecdotal case reports, vitamin D supplementation provided complete relief within a week in some patients having widespread, nonspecific pain that was unresponsive to analgesics, including opioids.<sup>79</sup> In other cases, pain and muscle weakness reportedly resolved “within

weeks” of beginning supplementation.<sup>72</sup>

In one study, pain relief from neuralgia was achieved at 3 months after beginning vitamin D supplementation.<sup>91</sup> In many cases, bone-related pain may require approximately 3 months of adequate vitamin D supplementation for its relief,<sup>47,65</sup> while muscle pain may need 6 months, and muscle weakness or fatigue may require even longer to resolve.<sup>47,65,76</sup>

Overall, Vasquez and colleagues<sup>37</sup> recommended that at least 5 to 9 months should be allowed for fully assessing either the benefits or ineffectiveness of vitamin D supplementation. Likewise, Vieth et al<sup>100</sup> suggested that the greatest physiologic responses may occur after 6 months of supplementation. Therefore, the timeframe recommended in Table 5 – monitoring results for up to 3 months – should be considered a minimum period of watchful waiting.

Complete pain relief would be easy for patients to detect, but in most cases this could be an unrealistic expectation. The evidence is suggestive of a potential range of improvements with vitamin D therapy, some more obvious than others. For example, instead of complete pain relief, patients may experience partial relief, reduced intensity or frequency of pain, less soreness or stiffness in muscles, increased stamina or strength, reductions in NSAID or opioid use, and/or improvements in mood or overall quality of life. These results are less spectacular than complete pain relief, but are still important and worthwhile outcomes to monitor.

## **Vitamin D Safety Considerations**

The highly favorable safety profile of vitamin D is evidenced by its lack of significant adverse effects, even at relatively high doses, and the absence of harmful interactions with other drugs. While vitamin D is potentially toxic, reports of associated overdoses and deaths have been relatively rare.

Tolerance & Toxicity. Excessive intake and accumulation of vitamin D is sometimes referred to as “hypervitaminosis D,” however this is poorly defined. Because a primary role of vitamin D is facilitating absorption of calcium from the intestine, the main signs/symptoms of vitamin D toxicity result from excessive serum calcium, or hypercalcemia (see Table 6).

The rather diverse signs/symptoms of hypercalcemia in patients with pain may be difficult to attribute to vitamin D intoxication, since they might mimic those of opioid side effects, neuropathy, or other conditions. Paradoxically, some symptoms match those of hypocalcemia. In some cases, patients with serum 25(OH)D at toxic levels can be clinically asymptomatic.<sup>126</sup>

Since full exposure to sunlight can provide the vitamin D<sub>3</sub> equivalent of up to 20,000 IU/day, the human body can obviously tolerate and safely manage relatively large daily doses. Toxicity has not been reported from repetitive daily exposure to sunlight.<sup>25</sup>

Still, supplementation via commercially manufactured vitamin D products would circumvent natural mechanisms in human skin that prevent excess D<sub>3</sub> production and accumulation resulting from sun exposure. A Tolerable Upper Intake Level, or UL, for oral vitamin D<sub>3</sub> supplementation—which is the long-term dose expected to pose no risk of observed adverse effects—currently is defined in the United States as 1000 IU/day in infants up to 12 months of age and 2000 IU/day for all other ages.<sup>16</sup> However, many experts assert that the 2000 IU/day UL is far too low.<sup>19,35-37,39,42</sup> An extensive review by Hathcock et al,<sup>121</sup> applying risk-assessment techniques, concluded that the UL for vitamin D consumption by adults actually could be 10,000 IU/day of D<sub>3</sub>, without risks of hypercalcemia.

However, the duration of high vitamin D intake may be the more critical factor. Taking 10,000 IU of D<sub>3</sub> daily for six months has been implicated as toxic,<sup>22,132</sup> whereas Holick [2007]<sup>14</sup> noted that this daily amount in adults can be well tolerated for five months. And, this 10,000 IU/day dose was used successfully and safely during three months in one study for relieving back pain.<sup>74</sup>

Vieth et al<sup>100</sup> demonstrated that 4000 IU of D3 per day was well tolerated during 15 months of therapy. Concentrations of 25(OH)D were increased to more optimal levels and parathyroid hormone was reduced, while calcium levels remained normal.

Some researchers have proposed that long-term daily consumption of 40,000 IU of vitamin D would be needed to cause hypercalcemia.<sup>18,19</sup> The US Office of Dietary Supplements<sup>133</sup> notes that hypercalcemia can result from 50,000 IU/day or more taken for an extended period of time.

There have not been any reports in the literature of a single, one-time excessive dose of vitamin D (D2 or D3) being toxic or fatal in humans. However, a number of incident reports of vitamin D toxicity involving very high doses, and including 6 fatalities, have appeared in the literature. In the incident reports, encompassing 77 cases of toxicity, amounts of vitamin D taken for periods ranging from days to years included daily doses from 160,000 IU up to an astounding 2.6 million IU.<sup>125,126,128,129,130,131,134,135</sup> The relatively few fatalities resulted from secondary causes during treatment for hypercalcemia. A common feature of all incidents was that victims were not knowingly or intentionally taking excessive amounts of vitamin D, and no one was taking vitamin D under practitioner supervision. In almost all cases, toxic overdoses could have been avoided with better quality control in product manufacture and/or education of patients in the proper use of supplements.

As another measure of safety, consolidated data for 2006 were examined (the most recently reported year) from 61 poison control centers serving 300 million persons in the United States.<sup>136</sup> There were only 516 mentions of incidents involving vitamin D which, by comparison, were roughly one-fourth the number for vitamin C and merely 0.8% of all incidents involving vitamin products. Only 13% of all vitamin D cases required treatment in a healthcare facility, although adverse signs/symptoms were absent or minimal in almost all patients (92%). Only 5 of the victims had more pronounced signs/symptoms of vitamin D toxicity, but these were not life threatening and there were no serious adverse events reported.

## **Vitamin D Drug Interactions**

There have been relatively few mentions in the literature of vitamin D supplements interacting with other agents or medications, and these are summarized in Table 7. In most cases, the potency of vitamin D is reduced by the other drug and the vitamin dose can be increased to accommodate this. Conversely, very high doses of vitamin D should be avoided due to possible risks of hypercalcemia when taken with digitalis/digoxin or certain diuretics.

Additionally, St. John's wort,<sup>27</sup> excessive alcohol,<sup>133</sup> and tobacco smoking<sup>138</sup> have been reported to potentially reduce the effects of vitamin D. Mineral oil and stimulant laxatives decrease dietary calcium absorption and can influence hypocalcemia.<sup>133</sup> Gastric bypass and other gastric or intestinal resection procedures have been associated with vitamin D insufficiency.<sup>60,106</sup>

It should be noted that not all patients would be affected by these interactions or effects, and vitamin D has not been noted to interfere harmfully with the actions of any medications. Therefore, none of the reported interactions or effects has been indicated in the literature as a contraindication for vitamin D supplementation.

## **Conclusions**

Extensive clinical evidence and expert commentary supports the opinion that recommending adequate vitamin D intake for helping patients with chronic musculoskeletal pain and fatigue syndromes should be more widely recognized and acted upon. In many cases, contributing factors are nonspecific or undetermined. Even in cases where a specific etiology has been diagnosed, the potential for vitamin D deficit as a factor contributing to and/or prolonging the pain condition should not be ruled out.

Table 6. Signs/Symptoms of D Toxicity (Hypercalcemia)

- abdominal pain
- achy muscles/joints
- anorexia
- azotemia
- calcifications
- constipation
- disorientation/confusion
- fatigue/lethargy
- fever/chills
- GI upset
- hypertension

References: Barrueto et al<sup>125</sup>; Blank et al<sup>126</sup>; Calcitrio<sup>112</sup>; Ca

Further clinical research studies would be helpful. Vitamin D is not proposed as a “cure” for all chronic pain conditions or in all patients. Optimal clinical outcomes of vitamin D therapy might be best attained via multicomponent treatment plans addressing many facets of health and pain relief. Therefore, vitamin D is not suggested as a replacement for any other approaches to pain management.

Table 7. Drugs Potentially Interacting with Vitamin D

- Potency of vitamin D is reduced. The vitamin dose ca
- antacids (aluminum- or magnesium-containing)
- anticonvulsants (eg, carbamazepine)
- antirejection meds (after organ transplant)
- antiretrovirals(HIV/AIDS therapies)
- barbiturates (eg, phenobarbital, phenytoin)
- cholestyramine
- colestipol
- corticosteroids
- glucocorticoids
- hydroxychloroquine
- rifampin

Risk of hypercalcemia when taken concurrently. Very high do

- digitalis/digoxin
- thiazide diuretics

References: Bringhurst et al<sup>137</sup>; Calcitriol<sup>12</sup>; Ergocalciferol<sup>1</sup>;  
Mascarenhas and Mobarhan<sup>51</sup>; ODS<sup>133</sup>; Turner et al<sup>60</sup>

To start, a conservative total daily supplementation of 2400 IU to 2800 IU of vitamin D3 is proposed as potentially benefitting patients. Along with that, some patience is advised regarding expectations for improvements; it may require up to 9 months before maximum effects are realized. In some cases, other factors or undetected conditions may be contributing to a chronic pain condition that vitamin D supplementation alone cannot ameliorate.

In sum, for patients with chronic musculoskeletal pain and related symptoms, supplemental vitamin D has a highly favorable benefit to cost ratio, with minimal, if any, risks. In all likelihood, it would do no harm and probably could do much good.n

## Acknowledgments

The author wishes to acknowledge and thank the following expert reviewers of the full research report

for their helpful comments and assistance: Bruce Hollis, PhD; Michael F. Holick, MD, PhD; Seth I. Kaufman, MD; Lee A. Kral, PharmD, BCPS; Paul W. Lofholm, PharmD, FACA; N. Lee Smith, MD; James D. Toombs, MD; Winnie Dawson, RN, BSN, MA.

## Disclosure

The author reports no involvement in the nutritional supplement industry or any other conflicts of interest. Covidien/ Mallinckrodt, St. Louis, MO, as a sponsor of Pain Treatment Topics, provided educational funding support for the research report on Vitamin D but had no role in its conception or development. The sponsor has no vested interests in the nutritional supplement industry and is not a manufacturer or distributor of vitamin products.

## References:

## Resources

- 1. Mohr SB, Garland CF, Gorham ED, Garland FC. The association between ultraviolet B irradiance, vitamin D status and incidence rates of type 1 diabetes in 51 regions worldwide. *Diabetologia*. 2008;51(8):1391-1398.
- 2. Ng K, Meyerhardt JA, Wu K, et al. Circulating 25-hydroxyvitamin D levels and survival in patients with colorectal cancer. *J Clin Oncol*. 2008. 26(18):2984-2991.
- 3. Giovannucci E, Liu Y, Hollis BW, Rimm EB. 25-hydroxyvitamin D and risk of myocardial infarction in men. *Arch Int Med*. 2008. 168(11):1174-1180.
- 4. APF (American Pain Foundation). Pain facts: overview of American pain surveys. 2007. Available online at: <http://www.painfoundation.org/page.asp?file=Newsroom/PainSurveys.htm>. Accessed 2/25/08.
- 5. Nachemson A, Waddell G, Norlund A. Epidemiology of neck and back pain. In: Nachemson A, Jonsson E (eds) *Neck and Back Pain: the Scientific Evidence of Causes, Diagnosis, and Treatment*. Phila, PA: Williams & Wilkins; 2000: 165-188.
- 6. NPF (National Pain Foundation). Harris Interactive Survey on Pain in America, Dec 2007. Released in 2008. Avail at: <http://www.nationalpainfoundation.org/MyResearch/PainSurvey2008.asp>. Accessed 2/25/08.
- 7. Watkins EA, Wollan PC, Melton III J, Yawn PB. A population in pain: report from the Olmsted County Health Study. *Pain Med*. 2008. 9(2):166-174.
- 8. Deyo RA. Diagnostic evaluation of LBP. *Arch Intern Med*. 2002. 162:1444-1448.
- 9. Deyo RA, Weinstein JN. Low back pain. *New Eng J Med*. 2001. 344:363-370.
- 10. Holick MF. Vitamin D for health and in chronic kidney disease. *Semin Dial*. 2005. 18:266-275.
- 11. Lau AH, How PP. The role of the pharmacist in the identification and management of secondary hyperparathyroidism. *US Pharmacist*. 2007(Jul):62-70. Avail at: <http://www.uspharmacist.com/index.asp?page=ce/105514/default.htm>. Accessed 2/15/08.
- 12. Calcitriol. Merck Manual [online]. 2007. Avail at: <http://www.merck.com/mmpe/lexicomp/calcitriol.html>. Accessed 2/19/08.
- 13. Ergocalciferol. Merck Manual [online]. 2007. Avail at: <http://www.merck.com/mmpe/lexicomp/ergocalciferol.html>. Accessed 2/19/08.
- 14. Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007. 357(3):266-281.
- 15. Marcus R. Agents affecting calcification and bone turnover: calcium, phosphate, parathyroid hormone, vitamin D, calcitonin, and other compounds. Chap 61. In: Hardman JG, Limbird LE, eds. *Goodman & Gilman's The Pharmacological Basis of Therapeutics*. 9th ed. New York, NY: McGraw-Hill; 1995.
- 16. ODS (Office of Dietary Supplements; US National Institutes of Health). Dietary Supplement Fact Sheet: Vitamin D. Updated Feb 19, 2008. Avail at: <http://dietary-supplements.info.nih.gov/factsheets/vitamind.asp>. Accessed 4/27/08
- 17. Singh YN. Vitamin D. Part I: Are we getting enough? *US Pharm*. 2004a. 10:66-72. Avail at: [http://www.uspharmacist.com/index.asp?show=article&page=8\\_1352.htm](http://www.uspharmacist.com/index.asp?show=article&page=8_1352.htm). Accessed 4/29/08.



- 18. Tavera-Mendoza LE, White JH. Cell defenses and the sunshine vitamin. *Scientific American*. Nov 2007. Avail at: <http://www.sciam.com/article.cfm?id=cell-defenses-and-the-sunshine-vitamin>. Accessed 3/1/08.
- 19. Vieth R. The pharmacology of vitamin D, including fortification strategies. Chapter 61. In: Feldman D, Glorieux F, eds. *Vitamin D*. 2nd ed. Boston, MA: Elsevier Academic Press; 2005. Avail at: <http://www.direct-ms.org/pdf/VitDVieth/Vieth%20chapter%2061.pdf>. Accessed 4/17/08.
- 20. Johnson LE. Vitamin D. *Merck Manual* [online]. 2007. Avail at: [http://www.merck.com/mmpe/sec01/ch004/ch004k.html?qt=vitamin%20D&alt=sh#tb004\\_6](http://www.merck.com/mmpe/sec01/ch004/ch004k.html?qt=vitamin%20D&alt=sh#tb004_6). Accessed 2/19/08.
- 21. Lips P. Vitamin D deficiency and secondary hyperparathyroidism in the elderly: consequences for bone loss and fractures and therapeutic implications. *Endocrine Rev*. 2001. 22(4):477-501.
- 22. Reginster J-Y. The high prevalence of inadequate serum vitamin D levels and implications for bone health. *Curr Med Res Opin*. 2005. 21(4):579-585. Avail at: [http://www.medscape.com/viewarticle/504783\\_print](http://www.medscape.com/viewarticle/504783_print). Accessed 2/24/08.
- 23. Holick MF, Biancuzzo RM, Chen TC, et al. Vitamin D<sub>2</sub> is as effective as vitamin D<sub>3</sub> in maintaining circulating concentrations of 25-hydroxyvitamin D. *J Clin Endocrinol Metab*. 2008. 93(3):677-687.
- 24. Vieth R. Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr*. 1999. 69:842-856.
- 25. Hollis BW. Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutr*. 2005. 135:317-322. Available by search at: <http://jn.nutrition.org/search.dtl>. Accessed 4/2/08.
- 26. Vieth R, Chan P-CR, MacFarlane GD. Efficacy and safety of vitamin D<sub>3</sub> intake exceeding the lowest observed adverse effect level. *Am J Clin Nutr*. 2001;73: 288-294. Avail at: <http://www.ajcn.org/cgi/content/full/73/2/288>.
- 27. Shinchuk L, Holick MF. Vitamin D and rehabilitation: improving functional outcomes. *Nutr Clin Prac*. 2007. 22(3):297-304.
- 28. Holick MF. Vitamin D: the underappreciated D-lightful hormone that is important for skeletal and cellular health. *Curr Opin Endocrinol Diabetes*. 2002b;9:87-98.
- 29. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem and health consequences. *Am J Clin Nutr*. 2008. 87(suppl):1080S-1086S.
- 30. Marshall TG. Vitamin D discovery outpaces FDA decision making. *Bioessays*. 2008. 30(2):173-182.
- 31. Walters MR. Newly identified actions of the vitamin D endocrine system. *Endocrine Rev*. 1992. 13(4):719-764.
- 32. Kawaguchi Y, Kanamori M, Ishihara H, et al. The association of lumbar disc disease and vitamin-D receptor gene polymorphism. *J Bone Joint Surg Am*. 2002. 84-A(11):2022-2028.
- 33. Videman T, Gibbons LE, Battie MC, et al. The relative roles of intragenic polymorphisms of the vitamin D receptor gene in lumbar spine degeneration and bone density. *Spine*. 2001. 26(3):A1-A6.
- 34. Grant WB. An estimate of premature cancer mortality in the U.S. due to inadequate doses of solar ultraviolet-B radiation. *Cancer*. 2002. 94:1867-1875.
- 35. Holick MF. Vitamin D: a millenium perspective. *J Cell Biochem*. 2003a. 88:296-307.
- 36. Holick MF. Vitamin D: photobiology, metabolism, mechanism of action, and clinical applications. In: Favus MJ, ed. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*. 5th ed. Washington, DC: American Society for Bone and Mineral Research. 2003c:129-137.
- 37. Vasquez A, Manso G, Cannell J. The clinical importance of vitamin D (cholecalciferol): a paradigm shift with implications for all healthcare providers. *Alternative Therapies*. 2004. 10(5):28-36. Avail at: <http://www.vitamindcouncil.com/PDFs/CME-clinicalImportanceVitD.pdf>. Accessed 2/21/08.
- 38. Heaney RP. Vitamin D: how much do we need, and how much is too much? *Osteoporosis Int*. 2000. 11:553-555.
- 39. Heaney RP. Functional indices of vitamin D status and ramifications of vitamin D deficiency. *Am J Clin Nutr*. 2004. 80(suppl):1706S-1709S.

- 40. Heaney RP, Dowell MS, Hale CA, Bendich A. Calcium absorption varies within the reference range for serum 25-hydroxyvitamin D. *J Am Coll Nutr.* 2003b. 22(2):142-146.
- 41. Reginster J-Y, Frederick I, Deroisy R, et al. Parathyroid hormone plasma concentrations in response to low 25-OH vitamin D levels increase with age in elderly women. *Osteoporosis Int.* 1998. 8:390-392.
- 42. Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr.* 2003a. 77:204-210.
- 43. Holick MF. Sunlight and vitamin D: both good for cardiovascular health [editorial]. *J Gen Intern Med.* 2002a. 17:733-735.
- 44. Holick MF. Vitamin D deficiency: what a pain it is. *Mayo Clin Proc.* 2003b. 78:1457-1459. Avail at: <http://www.mayoclinicproceedings.com/inside.asp?AID=464&UID>. Accessed 2/18/08.
- 45. Hickey L, Gordon CM. Vitamin D deficiency: new perspectives on an old disease. *Curr Opin Endocrinol Diabetes.* 2004. 11:18-25.
- 46. AHRQ (Agency for Healthcare Research and Quality). Cranney A, Horsley T, O'Donnell S, et al. Effectiveness and safety of vitamin D in relation to bone health. Evidence report/technology assessment No. 158 (prepared by the University of Ottawa Evidence-Based Practice Center [UO-EPC]). AHRQ publication 07-E013; Aug 2007. Avail at: <http://www.ahrq.gov/downloads/pub/evidence/pdf/vitamind/vitad.pdf>. Accessed 3/31/08.
- 47. Heath KM, Elovic EP. Vitamin D deficiency: implications in the rehabilitation setting. *Am J Phys Med Rehabil.* 2006. 85:916-923.
- 48. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc.* 2003. 78:1463-1470.
- 49. Jacobs ET, Alberts DS, Foote JA, et al. Vitamin D insufficiency in southern Arizona. *Am J Clin Nutr.* 2008. 87(3):608-613.
- 50. Gostine ML, Davis FN. Vitamin D deficiencies in pain patients. *Prac Pain Mgmt.* 2006(July/Aug):16-19.
- 51. Mascarenhas R, Mobarhan S. Hypovitaminosis D-induced pain. *Nutrition Reviews.* 2004. 62(9):354-359.
- 52. Bhattoa HP, Bettembuk P, Ganacharya S. Prevalence and seasonal variation of hypovitaminosis D and its relationship to bone metabolism in community dwelling postmenopausal Hungarian women. *Osteoporosis Int.* 2004. 15:447-451.
- 53. Hoogendijk VJG, Lips P, Dik MG, et al. Depression is associated with decreased 25-hydroxyvitamin D and increased parathyroid hormone in older adults. *Arch Gen Psychiatry.* 2008. 65(5):508-512.
- 54. Lotfi A, Abdel-Nasser AM, Hamdy A, Omran AA, El-Rehany MA. Hypovitaminosis D in female patients with chronic low back pain. *Clin Rheumatol.* 2007. 26:1895-1901.
- 55. MacFarlane GD, Sackrison JL Jr, Body JJ, et al. Hypovitaminosis D in a normal, apparently healthy urban European population. *J Steroid Biochem Mol Biol.* 2004. 89-90(1-5):621-622.
- 56. Nesby-O'Dell S, Scanlon KS, Cogswell ME, et al. Hypovitaminosis D prevalence and determinants among African American and white women of reproductive age: third National Health and Nutrition Examination Survey, 1988-1994. *Am J Clin Nutr.* 2002. 76:187-192.
- 57. Sullivan SS, Rosen CJ, Chen TC, Holick MF. Seasonal changes in serum 25(OH)D in adolescent girls in Maine [abstract]. *J Bone Miner Res.* 2003. 18(suppl 2):S407. Abstract M470.
- 58. Tangpricha V, Pearce EN, Chen TC, Holick MF. Vitamin D insufficiency among free-living healthy young adults. *Am J Med.* 2002. 112:659-662.
- 59. Yew KS, DeMieri PJ. Disorders of bone mineral metabolism. *Clin Fam Pract.* 2002. 4:522-572.
- 60. Turner MK, Hooten WM, Schmidt JE, et al. Prevalence and clinical correlates of vitamin D inadequacy among patients with chronic pain. *Pain Med.* 2008. online publication prior to print.
- 61. Hooten WM, Turner MK, Schmidt JE. Prevalence and clinical correlates of vitamin D in adequacy among patients with chronic pain. American Society of Anesthesiologists 2007 annual meeting, San Francisco. Oct 13-17, 2007. Abstract A1380.
- 62. Hicks GE, Shardell M, Miller RR, et al. Associations between vitamin D status and pain in older adults: the Invecchiare in Chianti Study. *J Am Geriatr Soc.* 2008. 56(5):785-791.

- 63. Armstrong DJ, Meenagh GK, Bickle I, et al. Vitamin D deficiency is associated with anxiety and depression in fibromyalgia [abstract]. Reported in: *J Musculoskeletal Pain*. 2007. 15(1):55.
- 64. Kealing J. Study: Joint pain ebbs with vitamin D. San Antonio Breast Cancer Symposium, Dec 16, 2007 - reported via *LJWorld.com*. Avail at: [http://www2.ljworld.com/news/2007/dec/16/study\\_joint\\_pain\\_ebbs\\_vitamin\\_d/](http://www2.ljworld.com/news/2007/dec/16/study_joint_pain_ebbs_vitamin_d/). Accessed 5/2/08.
- 65. de la Jara GDT, Pecoud A, Favrat B. Female asylum seekers with musculoskeletal pain: the importance of diagnosis and treatment of hypovitaminosis D. *BMC Family Practice*. 2006. 7(4). Avail at: <http://www.biomedcentral.com/1471-2296/7/4>. Accessed 3/8/08
- 66. Benson J, Wilson A, Stocks N, Moulding N. Muscle pain as an indicator of vitamin D deficiency in an urban Australian Aboriginal population. *Med J Australia*. 2006. 185(2):76-77. Avail at: [http://www.mja.com.au/public/issues/185\\_02\\_170706/ben10209\\_fm.html](http://www.mja.com.au/public/issues/185_02_170706/ben10209_fm.html). Access checked 3/3/08
- 67. Helliwell PS, Ibrahim GH, Karim Z, Sokoll K, Johnson H. Unexplained musculoskeletal pain in people of South Asian ethnic group referred to a rheumatology clinic - relationship to biochemical osteomalacia, persistence over time and response to treatment with calcium and vitamin D. *Clin Exp Rheumatol*. 2006. 24(4):424-427.
- 68. Erkal MZ, Wilde J, Bilgin Y, et al. High prevalence of vitamin D deficiencies, secondary to hyper-parathyroidism and generalized bone pain in Turkish immigrants in Germany: identification of risk factors. *Osteoporosis Int*. 2006. 17(8):1133-1140.
- 69. Macfarlane GJ, Palmer B, Roy D, et al. An excess of widespread pain among South Asians: are low levels of vitamin D implicated? *Ann Rheum Dis*. 2005. 64(8):1217-1219.
- 70. Baker K, Zhang YQ, Goggins J, et al. Hypovitaminosis D and its association with muscle strength, pain and physical function in knee osteoarthritis (OA): a 30-month longitudinal, observational study. American College of Rheumatology meeting; San Antonio, TX; Oct 16-21, 2004. Abstract 1755.
- 71. Haque U, Regan MJ, Nanda S. et al. Vitamin D: are we ignoring an important marker in patients with chronic musculoskeletal symptoms? American College of Rheumatology meeting; San Antonio, TX; Oct 16-21, 2004. Abstract 694.
- 72. van der Heyden JJ, Verrips A, ter Laak HJ, et al. Hypovitaminosis D-related myopathy in immigrant teenagers. *Neuropediatrics*. 2004. 35(5):290-292.
- 73. Block SR. Vitamin D deficiency is not associated with nonspecific musculoskeletal pain syndromes including fibromyalgia [letter]. *Mayo Clin Proc*. 2004. 79:1585-1591. Avail at: <http://www.mayoclinicproceedings.com/inside.asp?AID=2257&UID>. Access checked 3/12/08.
- 74. Al Faraj S, Al Mutairi K. Vitamin D deficiency and chronic low back pain in Saudi Arabia. *Spine* 2003. 28:177-179.
- 75. Huisman AM, White KP, Algra A, et al. Vitamin D levels in women with systematic lupus erythematosus and fibromyalgia. *J Rheumatol*. 2001. 28(11):2535-2539.
- 76. Glerup H, Mikkelsen K, Poulsen L, et al. Hypovitaminosis D myopathy without biochemical signs of osteomalacic bone involvement. *Calcif Tissue Int*. 2000b. 66:419-424.
- 77. Prabhala A, Garg R, Dandona P. Severe myopathy associated with vitamin D deficiency in western New York. *Arch Intern Med*. 2000. 160(8):1199-1203.
- 78. McAlindon TE, Felson DT, Zhang Y, et al. Relation of dietary intake and serum levels of vitamin D to progression of osteoarthritis of the knee among participants in the Framingham Study. *Ann Int Med*. 1996. 125(5):353-359. Avail at: <http://www.annals.org/cgi/content/full/125/5/353>. Accessed 4/21/08
- 79. Gloth FM 3rd, Lindsay JM, Zelesnick LB, Greenough WB 3rd. Can vitamin D deficiency produce an unusual pain syndrome? *Arch Intern Med*. 1991. 151:1662-1664.
- 80. Boland R. Role of vitamin D in skeletal muscle function. *Endocr Rev*. 1986;7:434-438.
- 81. Haddad JG, Walgate J, Min C, Hahn TJ. Vitamin D metabolite-binding proteins in human tissue. *Biochem Biophys Acta*. 1976;444:921-925.
- 82. Schott G, Wills M. Muscle weakness in osteomalacia. *Lancet*. 1976;2:626-629.
- 83. Simpson RU, Thomas GA, Arnold AJ. Identification of 1,25-dihydroxyvitamin D3 receptors and activities in muscle. *J Biol Chem*. 1985. 260:8882-8891.
- 84. Pfeifer M, Begerow B, Minne HW, et al. Vitamin D status, trunk muscle strength, body sway, falls, and fractures among 237 postmenopausal women with osteoporosis. *Exp Clin Endocrinol*

Diabetes. 2001. 109:87-92.

- 85. Bischoff HA, Stahelin HB, Dick W, et al. Effects of vitamin D and calcium supplementation on falls: a randomized controlled trial. *J Bone Miner Res*. 2003. 18:343-351.
- 86. Bischoff-Ferrari HA, Willett WC, Wong JB, et al. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA*. 2005. 293(18):2257-2264.
- 87. Bischoff-Ferrari HA, Giovannucci E, Willett WC, et al. Estimation of optimum serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr*. 2006. 84:18-28.
- 88. Dukas L, Bischoff HA, Lindpainter LS, et al. Alfacalcidol reduces the number of fallers in a community-dwelling elderly population with a minimum calcium intake of more than 500 mg daily. *J Am Geriatr Soc*. 2004. 52:230-236.
- 89. Sambrook PN, Chen JS, March LM, et al. Serum parathyroid hormone predicts time to fall independent of vitamin D status in a frail elderly population. *J Clin Endocrinol Metab*. 2004. 89:1572-1576.
- 90. Boxer RS, Dauser RA, Walsh SJ, et al. The association between vitamin D and inflammation with the 6-minute walk and frailty in patients with heart failure. *J Am Geriatr Soc*. 2008. 56:454-461.
- 91. Lee P, Chen R. Vitamin D as an analgesic for patients with type 2 diabetes and neuropathic pain. *Arch Intern Med*. 2008. 168(7):771-772.
- 92. D'Ambrosio D, Cippitelli M, Cocciolo MG, et al. Inhibition of IL-12 production by 1,25-Dihydroxyvitamin D<sub>3</sub>. *J Clin Invest*. 1998. 101:252-262.
- 93. Schleithoff SS, Zittermann A, Tenderich G, et al. Vitamin D supplementation improves cytokine profiles in patients with congestive heart failure: a double-blind randomized, placebo-controlled trial. *Am J Clin Nutr*. 2006. 83:754-759.
- 94. Van den Berghe, Van Roosbroeck D, Vanhove P, et al. Bone turnover in prolonged critical illness: effect of vitamin D. *J Clin Endocr Metab*. 2003. 88:4623-4632.
- 95. Thys-Jacobs S. Alleviation of migraine with therapeutic vitamin D and calcium. *Headache*. 1994a. 34(10):590-592.
- 96. Thys-Jacobs S. Vitamin D and calcium in menstrual migraine. *Headache*. 1994b;34(9):544-546.
- 97. Gloth FM 3rd, Alam W, Hollis B. Vitamin D vs broad spectrum phototherapy in the treatment of seasonal affective disorder. *J Nutr Health Aging*. 1999. 3(1):5-7.
- 98. Lansdowne AT, Provost SC. Vitamin D<sub>3</sub> enhances mood in healthy subjects during winter. *Psychopharmacology*. 1998. 135(4):319-323.
- 99. Basaran S, Guzel R, Coskun-Benliday I, Guler-Uysal F. Vitamin D status: effects on quality of life in osteoporosis among Turkish women. *Qaul Life Res*. 2007. 16(9):1491-1499.
- 100. Vieth R, Kimball S, Hu A, Walfish PG. Randomized comparison of the effects of the vitamin D<sub>3</sub> adequate intake versus 100mcg (4000 IU) per day on biochemical responses and the wellbeing of patients. *Nutrition J*. 2004. 3:8. Avail at: <http://www.nutritionj.com/content/pdf/1475-2891-3-8.pdf>. Accessed 5/14/08.
- 101. Smith GR, Collinson PO, Kiely PD. Diagnosing hypovitaminosis D: serum measurements of calcium, phosphate, and alkaline phosphatase are unreliable, even in the presence of secondary hyper-parathyroidism. *Rheumatol*. 2005. 32(4):684-689.
- 102. Nellen JFJB, Smulders YM, PH Jos Frissen, EH Slaats, J Silberbusch. Hypovitaminosis D in immigrant women: slow to be diagnosed. *BMJ*. 1996. 312:570-572.
- 103. Peach H, Compston JE, Vedi S, Horton LWL. Value of plasma calcium, phosphate, and alkaline phosphate measurements in the diagnosis of histological osteomalacia. *J Clin Pathol*. 1982;35:625-630. Avail at: <http://jcp.bmj.com/cgi/reprint/35/6/625>. Accessed 3/7/08.
- 104. Cooper MS, Gittoes NJL. Diagnosis and management of hypocalcemia. *BMJ*. 2008. 336:1298-1302.
- 105. Fitzpatrick L. Hypocalcemia: Diagnosis and Treatment - Chapter 7. 2002. Endotext.com. Avail at:
  - <http://www.endotext.org/parathyroid/parathyroid7/parathyroid7.htm>. Accessed 2/23/08.
- 106. Lyman D. Undiagnosed vitamin D deficiency in the hospitalized patient. *Am Fam Phys*.

2005. 71(2):299-304. Avail at: <http://www.aafp.org/afp/20050115/299.pdf>. Accessed 4/30/08.
- 107. Skugor M, Milas M. Hypocalcemia. 2004. Cleveland Clinic; Avail at: <http://www.clevelandclinicmeded.com/>. Accessed 2/23/08.
  - 108. Peacey SR. Routine biochemistry in suspected vitamin D deficiency. *JR Soc Med*. 2004. 97:322-325. Avail at: <http://www.jrsm.org/cgi/reprint/97/7/322>. Accessed 3/8/08.
  - 109. Zerwekh JE. Blood biomarkers of vitamin D status. *Am J Clin Nutr*. 2008. 87(suppl):1087S-1094S.
  - 110. Singh RJ. Are clinical laboratories prepared for accurate testing of 25-hydroxy vitamin D? [letter]. *Clin Chem*. 2008. 54(1):221-223.
  - 111. Masood H, Narang AP, Bhat IA, Shah GN. Persistent limb pain and raised serum alkaline phosphatase the earliest markers to subclinical hypovitaminosis D in Kashmir [abstract]. *Indian J Physiol Pharmacol*. 1989. 33(4):259-261.
  - 112. Holick MF. Vitamin D deficiency as a contributor to multiple forms of chronic pain - reply II [letter]. *Mayo Clin Proc*. 2004a;79:694-709. Avail at: <http://www.mayoclinicproceedings.com/inside.asp?AID=2197&UID>. Accessed 3/7/08.
  - 113. IOM (Institute of Medicine). Standing Committee on the Scientific Evaluation of Dietary Reference Intakes. Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride. Washington, DC: National Academy Press. 1997. 71-145.
  - 114. DHHS (US Department of Health and Human Services and US Dept of Agriculture). Dietary Guidelines for Americans, 2005. 6th ed. Washington, DC: US Government Printing Office. 2005(Jan). Avail at: <http://www.health.gov/dietaryguidelines/dga2005/document/>. Accessed 5/18/08.
  - 115. Armas LAG, Hollis BW, Heaney RP. Vitamin D2 is much less effective than vitamin D3 in humans. *J Clin Endocrinol Metab*. 2004. 89(11):5387-5391. Avail at: <http://jcem.endojournals.org/cgi/reprint/89/11/5387>. Accessed 4/23/08.
  - 116. Barger-Lux MJ, Heaney RP, Dowell S, Chen TC, Holick MF. Vitamin D and its major metabolites: serum levels after graded oral dosing in healthy men. *Osteoporos Int*. 1998. 8(3):222-230.
  - 117. Cranney A, Horsley T, O'Donnell S, et al. Effectiveness and safety of vitamin D in relation to bone health. *Evid Rep Technol Assess*. 2007. 158:1-235.
  - 118. Trang HM, Cole DEC, Rubin LA, et al. Evidence that vitamin D3 increases serum 25-hydroxyvitamin D more efficiently than does vitamin D2. *Am J Clin Nutr*. 1998. 68:854-858. Avail at: <http://www.ajcn.org/cgi/reprint/68/4/854>. Accessed 4/23/08.
  - 119. Vieth R, Bischoff-Ferrari H, Boucher BJ. The urgent need to recommend an intake of vitamin D that is effective. *Am J Clin Nutr*. 2007. 85:649-650.
  - 120. Smith H, Anderson F, Raphael P, et al. Effect of annual intramuscular vitamin D on fracture risk in elderly men and women - a population-based, randomized, double-blind, placebo-controlled trial. *Rheumatology*. 2007. 46(12):1852-1857.
  - 121. Hathcock JN, Shao A, Vieth R, Heaney R. Risk assessment for vitamin D. *Am J Clin Nutr*. 2007. 85:6-18.
  - 122. Trivedi DP, Doll R, Khaw KT. Effect of four monthly oral vitamin D3 (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomized double blind controlled trial. *Br Med J*. 2003. 326:469.
  - 123. Romagnoli E, Mascia ML, Cipriani C, et al. Short and long term variations in serum calcitropic hormones after a single very large dose of ergocalciferol (vitamin D2) or cholecalciferol (vitamin D3) in the elderly. *J Clin Endocrinol Metab*. 2008. online publication ahead of print; doi:10.1210/jc.2008-0350.
  - 124. Hollis BW, MacFarlane G, Valcour A. Clinical significance and assessment of 25 OH vitamin D. Burlington, NC: LabCorp; 2005. Avail at: [http://www.labcorp.com/cme/pdf/Vitamin\\_D\\_Journal\\_Article.pdf](http://www.labcorp.com/cme/pdf/Vitamin_D_Journal_Article.pdf). Accessed 4/16/08.
  - 125. Barrueto F Jr, Wang-Flores HH, Howland MA, Hoffman RS, Nelson LS. Acute vitamin D intoxication in a child. *Pediatrics*. 2005. 116:e453-e456.
  - 126. Blank S, Scanlon KS, Sinks TH, Lett S, Falk H. An outbreak of hypervitaminosis D associated with the over fortification of milk from a home-delivery dairy. *Am J Pub Health*. 1995. 85(5):656-659.

- 127. Carroll MF, Schade DS. A practical approach to hypercalcemia. *Am Fam Physician*. 2003. 67:1959-1966. Avail at: <http://www.aafp.org/afp/20030501/1959.html>. Accessed 4/21/08.
- 128. Klontz KC, Acheson DW. Dietary supplement-induced vitamin D intoxication [letter]. *New Eng J Med*. 2007. 357:308-309.
- 129. Koutkia P, Chen TC, Holick MF. Vitamin D intoxication associated with an over-the-counter supplement. *New Engl J Med*. 2001. 345(1):65-66.
- 130. Todd MA, Bailey RR, Espiner EA, Lynn KL. Vitamin D2 for the treatment of chilblains – a cautionary tale. *N Z Med J*. 1987;100:465.
- 131. Vieth R, Pinto TR, Reen BS, Wong MM. Vitamin D poisoning by table sugar. *Lancet*. 2002. 359:672.
- 132. Jacobus CH, Holick MF, Shao Q, et al. Hypervitaminosis D associated with drinking milk. *New Engl J Med*. 1992;326:1173-1177.
- 133. ODS (Office of Dietary Supplements; US National Institutes of Health). Dietary Supplement Fact Sheet: Calcium. Updated Sept 23, 2005. Avail at: [http://dietary-supplements.info.nih.gov/factsheets/Calcium\\_pf.asp](http://dietary-supplements.info.nih.gov/factsheets/Calcium_pf.asp). Accessed 4/16/08
- 134. Adams JS, Lee G. Gains in bone mineral density with resolution of vitamin D intoxication. *Ann Int Med*. 1997. 127(3):203-206.
- 135. Pettifor JM, Bikle DD, Cavaleros M. Serum levels of free 1,25-dihydroxyvitamin D in vitamin D toxicity. *Ann Int Med*. 1995. 122(7):511-513.
- 136. Bronstein AC, Spyker DA, Cantilene Jr LR, et al. 2006 Annual Report of the American Association of Poison Control Centers National Poison Data System (NPDS). *Clinical Toxicology*. 2007. 45(8):815-917. Avail at: <http://www.aapcc.org/poison1.htm>. Accessed 2/22/08.
- 137. Bringhurst RF, Bemay MB, Krune SM, Kronenberg HM. Bone and mineral metabolism in health and disease. In: Kasper DL, et al. eds. *Harrison's Principles of Internal Medicine*. 16th Ed. New York: McGraw-Hill; 2005: 2238-2249.

[View Sources](#) [4]

## Resources

- 1. Mohr SB, Garland CF, Gorham ED, Garland FC. The association between ultraviolet B irradiance, vitamin D status and incidence rates of type 1 diabetes in 51 regions worldwide. *Diabetologia*. 2008;51(8):1391-1398.
- 2. Ng K, Meyerhardt JA, Wu K, et al. Circulating 25-hydroxyvitamin D levels and survival in patients with colorectal cancer. *J Clin Oncol*. 2008. 26(18):2984-2991.
- 3. Giovannucci E, Liu Y, Hollis BW, Rimm EB. 25-hydroxyvitamin D and risk of myocardial infarction in men. *Arch Int Med*. 2008. 168(11):1174-1180.
- 4. APF (American Pain Foundation). Pain facts: overview of American pain surveys. 2007. Available online at: <http://www.painfoundation.org/page.asp?file=Newsroom/PainSurveys.htm>. Accessed 2/25/08.
- 5. Nachemson A, Waddell G, Norlund A. Epidemiology of neck and back pain. In: Nachemson A, Jonsson E (eds) *Neck and Back Pain: the Scientific Evidence of Causes, Diagnosis, and Treatment*. Phila, PA: Williams & Wilkins; 2000: 165-188.
- 6. NPF (National Pain Foundation). Harris Interactive Survey on Pain in America, Dec 2007. Released in 2008. Avail at: <http://www.nationalpainfoundation.org/MyResearch/PainSurvey2008.asp>. Accessed 2/25/08.
- 7. Watkins EA, Wollan PC, Melton III J, Yawn PB. A population in pain: report from the Olmsted County Health Study. *Pain Med*. 2008. 9(2):166-174.
- 8. Deyo RA. Diagnostic evaluation of LBP. *Arch Intern Med*. 2002. 162:1444-1448.
- 9. Deyo RA, Weinstein JN. Low back pain. *New Eng J Med*. 2001. 344:363-370.
- 10. Holick MF. Vitamin D for health and in chronic kidney disease. *Semin Dial*. 2005. 18:266-275.
- 11. Lau AH, How PP. The role of the pharmacist in the identification and management of secondary hyperparathyroidism. *US Pharmacist*. 2007(Jul):62-70. Avail at:

<http://www.uspharmacist.com/index.asp?page=ce/105514/default.htm>. Accessed 2/15/08.

- 12. Calcitriol. Merck Manual [online]. 2007. Avail at: <http://www.merck.com/mmpe/lexicomp/calcitriol.html>. Accessed 2/19/08.
- 13. Ergocalciferol. Merck Manual [online]. 2007. Avail at: <http://www.merck.com/mmpe/lexicomp/ergocalciferol.html>. Accessed 2/19/08.
- 14. Holick MF. Vitamin D deficiency. *N Engl J Med*. 2007. 357(3):266-281.
- 15. Marcus R. Agents affecting calcification and bone turnover: calcium, phosphate, parathyroid hormone, vitamin D, calcitonin, and other compounds. Chap 61. In: Hardman JG, Limbird LE, eds. *Goodman & Gilman's The Pharmacological Basis of Therapeutics*. 9th ed. New York, NY: McGraw-Hill; 1995.
- 16. ODS (Office of Dietary Supplements; US National Institutes of Health). Dietary Supplement Fact Sheet: Vitamin D. Updated Feb 19, 2008. Avail at: <http://dietary-supplements.info.nih.gov/factsheets/vitaminD.asp>. Accessed 4/27/08
- 17. Singh YN. Vitamin D. Part I: Are we getting enough? *US Pharm*. 2004a. 10:66-72. Avail at: [http://www.uspharmacist.com/index.asp?show=article&page=8\\_1352.htm](http://www.uspharmacist.com/index.asp?show=article&page=8_1352.htm). Accessed 4/29/08.
- 18. Tavera-Mendoza LE, White JH. Cell defenses and the sunshine vitamin. *Scientific American*. Nov 2007. Avail at: <http://www.sciam.com/article.cfm?id=cell-defenses-and-the-sunshine-vitamin>. Accessed 3/1/08.
- 19. Vieth R. The pharmacology of vitamin D, including fortification strategies. Chapter 61. In: Feldman D, Glorieux F, eds. *Vitamin D*. 2nd ed. Boston, MA: Elsevier Academic Press; 2005. Avail at: <http://www.direct-ms.org/pdf/VitDVieth/Vieth%20chapter%2061.pdf>. Accessed 4/17/08.
- 20. Johnson LE. Vitamin D. Merck Manual [online]. 2007. Avail at: [http://www.merck.com/mmpe/sec01/ch004/ch004k.html?qt=vitamin%20D&alt=sh#tb004\\_6](http://www.merck.com/mmpe/sec01/ch004/ch004k.html?qt=vitamin%20D&alt=sh#tb004_6). Accessed 2/19/08.
- 21. Lips P. Vitamin D deficiency and secondary hyperparathyroidism in the elderly: consequences for bone loss and fractures and therapeutic implications. *Endocrine Rev*. 2001. 22(4):477-501.
- 22. Reginster J-Y. The high prevalence of inadequate serum vitamin D levels and implications for bone health. *Curr Med Res Opin*. 2005. 21(4):579-585. Avail at: [http://www.medscape.com/viewarticle/504783\\_print](http://www.medscape.com/viewarticle/504783_print). Accessed 2/24/08.
- 23. Holick MF, Biancuzzo RM, Chen TC, et al. Vitamin D<sub>2</sub> is as effective as vitamin D<sub>3</sub> in maintaining circulating concentrations of 25-hydroxyvitamin D. *J Clin Endocrinol Metab*. 2008. 93(3):677-687.
- 24. Vieth R. Vitamin D supplementation, 25-hydroxyvitamin D concentrations, and safety. *Am J Clin Nutr*. 1999. 69:842-856.
- 25. Hollis BW. Circulating 25-hydroxyvitamin D levels indicative of vitamin D sufficiency: implications for establishing a new effective dietary intake recommendation for vitamin D. *J Nutr*. 2005. 135:317-322. Available by search at: <http://jn.nutrition.org/search.dtl>. Accessed 4/2/08.
- 26. Vieth R, Chan P-CR, MacFarlane GD. Efficacy and safety of vitamin D<sub>3</sub> intake exceeding the lowest observed adverse effect level. *Am J Clin Nutr*. 2001;73: 288-294. Avail at: <http://www.ajcn.org/cgi/content/full/73/2/288>.
- 27. Shinchuk L, Holick MF. Vitamin D and rehabilitation: improving functional outcomes. *Nutr Clin Prac*. 2007. 22(3):297-304.
- 28. Holick MF. Vitamin D: the underappreciated D-lightful hormone that is important for skeletal and cellular health. *Curr Opin Endocrinol Diabetes*. 2002b;9:87-98.
- 29. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem and health consequences. *Am J Clin Nutr*. 2008. 87(suppl):1080S-1086S.
- 30. Marshall TG. Vitamin D discovery outpaces FDA decision making. *Bioessays*. 2008. 30(2):173-182.
- 31. Walters MR. Newly identified actions of the vitamin D endocrine system. *Endocrine Rev*. 1992. 13(4):719-764.
- 32. Kawaguchi Y, Kanamori M, Ishihara H, et al. The association of lumbar disc disease and vitamin-D receptor gene polymorphism. *J Bone Joint Surg Am*. 2002. 84-A(11):2022-2028.
- 33. Videman T, Gibbons LE, Battie MC, et al. The relative roles of intragenic polymorphisms of the vitamin D receptor gene in lumbar spine degeneration and bone density. *Spine*. 2001. 26(3):A1-A6.

- 34. Grant WB. An estimate of premature cancer mortality in the U.S. due to inadequate doses of solar ultraviolet-B radiation. *Cancer*. 2002. 94:1867-1875.
- 35. Holick MF. Vitamin D: a millenium perspective. *J Cell Biochem*. 2003a. 88:296-307.
- 36. Holick MF. Vitamin D: photobiology, metabolism, mechanism of action, and clinical applications. In: Favus MJ, ed. *Primer on the Metabolic Bone Diseases and Disorders of Mineral Metabolism*. 5th ed. Washington, DC: American Society for Bone and Mineral Research. 2003c:129-137.
- 37. Vasquez A, Manso G, Cannell J. The clinical importance of vitamin D (cholecalciferol): a paradigm shift with implications for all healthcare providers. *Alternative Therapies*. 2004. 10(5):28-36. Avail at:
  - <http://www.vitamindcouncil.com/PDFs/CME-clinicalImportanceVitD.pdf>. Accessed 2/21/08.
- 38. Heaney RP. Vitamin D: how much do we need, and how much is too much? *Osteoporosis Int*. 2000. 11:553-555.
- 39. Heaney RP. Functional indices of vitamin D status and ramifications of vitamin D deficiency. *Am J Clin Nutr*. 2004. 80(suppl):1706S-1709S.
- 40. Heaney RP, Dowell MS, Hale CA, Bendich A. Calcium absorption varies within the reference range for serum 25-hydroxyvitamin D. *J Am Coll Nutr*. 2003b. 22(2):142-146.
- 41. Reginster J-Y, Frederick I, Deroisy R, et al. Parathyroid hormone plasma concentrations in response to low 25-OH vitamin D levels increase with age in elderly women. *Osteoporosis Int*. 1998. 8:390-392.
- 42. Heaney RP, Davies KM, Chen TC, Holick MF, Barger-Lux MJ. Human serum 25-hydroxycholecalciferol response to extended oral dosing with cholecalciferol. *Am J Clin Nutr*. 2003a. 77:204-210.
- 43. Holick MF. Sunlight and vitamin D: both good for cardiovascular health [editorial]. *J Gen Intern Med*. 2002a. 17:733-735.
- 44. Holick MF. Vitamin D deficiency: what a pain it is. *Mayo Clin Proc*. 2003b. 78:1457-1459. Avail at: <http://www.mayoclinicproceedings.com/inside.asp?AID=464&UID>. Accessed 2/18/08.
- 45. Hickey L, Gordon CM. Vitamin D deficiency: new perspectives on an old disease. *Curr Opin Endocrinol Diabetes*. 2004. 11:18-25.
- 46. AHRQ (Agency for Healthcare Research and Quality). Cranney A, Horsley T, O'Donnell S, et al. Effectiveness and safety of vitamin D in relation to bone health. Evidence report/technology assessment No. 158 (prepared by the University of Ottawa Evidence-Based Practice Center [UO-EPC]). AHRQ publication 07-E013; Aug 2007. Avail at: <http://www.ahrq.gov/downloads/pub/evidence/pdf/vitamind/vitad.pdf>. Accessed 3/31/08.
- 47. Heath KM, Elovic EP. Vitamin D deficiency: implications in the rehabilitation setting. *Am J Phys Med Rehabil*. 2006. 85:916-923.
- 48. Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent, nonspecific musculoskeletal pain. *Mayo Clin Proc*. 2003. 78:1463-1470.
- 49. Jacobs ET, Alberts DS, Foote JA, et al. Vitamin D insufficiency in southern Arizona. *Am J Clin Nutr*. 2008. 87(3):608-613.
- 50. Gostine ML, Davis FN. Vitamin D deficiencies in pain patients. *Prac Pain Mgmt*. 2006(July/Aug):16-19.
- 51. Mascarenhas R, Mobarhan S. Hypovitaminosis D-induced pain. *Nutrition Reviews*. 2004. 62(9):354-359.
- 52. Bhattoa HP, Bettembuk P, Ganacharya S. Prevalence and seasonal variation of hypovitaminosis D and its relationship to bone metabolism in community dwelling postmenopausal Hungarian women. *Osteoporosis Int*. 2004. 15:447-451.
- 53. Hoogendijk VJG, Lips P, Dik MG, et al. Depression is associated with decreased 25-hydroxyvitamin D and increased parathyroid hormone in older adults. *Arch Gen Psychiatry*. 2008. 65(5):508-512.
- 54. Lotfi A, Abdel-Nasser AM, Hamdy A, Omran AA, El-Rehany MA. Hypovitaminosis D in female patients with chronic low back pain. *Clin Rheumatol*. 2007. 26:1895-1901.
- 55. MacFarlane GD, Sackrison JL Jr, Body JJ, et al. Hypovitaminosis D in a normal, apparently healthy urban European population. *J Steroid Biochem Mol Biol*. 2004. 89-90(1-5):621-622.
- 56. Nesby-O'Dell S, Scanlon KS, Cogswell ME, et al. Hypovitaminosis D prevalence and



determinants among African American and white women of reproductive age: third National Health and Nutrition Examination Survey, 1988-1994. *Am J Clin Nutr.* 2002. 76:187-192.

- 57. Sullivan SS, Rosen CJ, Chen TC, Holick MF. Seasonal changes in serum 25(OH)D in adolescent girls in Maine [abstract]. *J Bone Miner Res.* 2003. 18(suppl 2):S407. Abstract M470.
- 58. Tangpricha V, Pearce EN, Chen TC, Holick MF. Vitamin D insufficiency among free-living healthy young adults. *Am J Med.* 2002. 112:659-662.
- 59. Yew KS, DeMieri PJ. Disorders of bone mineral metabolism. *Clin Fam Pract.* 2002. 4:522-572.
- 60. Turner MK, Hooten WM, Schmidt JE, et al. Prevalence and clinical correlates of vitamin D inadequacy among patients with chronic pain. *Pain Med.* 2008. online publication prior to print.
- 61. Hooten WM, Turner MK, Schmidt JE. Prevalence and clinical correlates of vitamin D in adequacy among patients with chronic pain. American Society of Anesthesiologists 2007 annual meeting, San Francisco. Oct 13-17, 2007. Abstract A1380.
- 62. Hicks GE, Shardell M, Miller RR, et al. Associations between vitamin D status and pain in older adults: the Invecchiare in Chianti Study. *J Am Geriatr Soc.* 2008. 56(5):785-791.
- 63. Armstrong DJ, Meenagh GK, Bickle I, et al. Vitamin D deficiency is associated with anxiety and depression in fibromyalgia [abstract]. Reported in: *J Musculoskeletal Pain.* 2007. 15(1):55.
- 64. Kealing J. Study: Joint pain ebbs with vitamin D. San Antonio Breast Cancer Symposium, Dec 16, 2007 - reported via *LJWorld.com*. Avail at: [http://www2.ljworld.com/news/2007/dec/16/study\\_joint\\_pain\\_ebbs\\_vitamin\\_d/](http://www2.ljworld.com/news/2007/dec/16/study_joint_pain_ebbs_vitamin_d/). Accessed 5/2/08.
- 65. de la Jara GDT, Pecoud A, Favrat B. Female asylum seekers with musculoskeletal pain: the importance of diagnosis and treatment of hypovitaminosis D. *BMC Family Practice.* 2006. 7(4). Avail at: <http://www.biomedcentral.com/1471-2296/7/4>. Accessed 3/8/08
- 66. Benson J, Wilson A, Stocks N, Moulding N. Muscle pain as an indicator of vitamin D deficiency in an urban Australian Aboriginal population. *Med J Australia.* 2006. 185(2):76-77. Avail at: [http://www.mja.com.au/public/issues/185\\_02\\_170706/ben10209\\_fm.html](http://www.mja.com.au/public/issues/185_02_170706/ben10209_fm.html). Access checked 3/3/08
- 67. Helliwell PS, Ibrahim GH, Karim Z, Sokoll K, Johnson H. Unexplained musculoskeletal pain in people of South Asian ethnic group referred to a rheumatology clinic - relationship to biochemical osteomalacia, persistence over time and response to treatment with calcium and vitamin D. *Clin Exp Rheumatol.* 2006. 24(4):424-427.
- 68. Erkal MZ, Wilde J, Bilgin Y, et al. High prevalence of vitamin D deficiencies, secondary to hyper-parathyroidism and generalized bone pain in Turkish immigrants in Germany: identification of risk factors. *Osteoporosis Int.* 2006. 17(8):1133-1140.
- 69. Macfarlane GJ, Palmer B, Roy D, et al. An excess of widespread pain among South Asians: are low levels of vitamin D implicated? *Ann Rheum Dis.* 2005. 64(8):1217-1219.
- 70. Baker K, Zhang YQ, Goggins J, et al. Hypovitaminosis D and its association with muscle strength, pain and physical function in knee osteoarthritis (OA): a 30-month longitudinal, observational study. American College of Rheumatology meeting; San Antonio, TX; Oct 16-21, 2004. Abstract 1755.
- 71. Haque U, Regan MJ, Nanda S. et al. Vitamin D: are we ignoring an important marker in patients with chronic musculoskeletal symptoms? American College of Rheumatology meeting; San Antonio, TX; Oct 16-21, 2004. Abstract 694.
- 72. van der Heyden JJ, Verrips A, ter Laak HJ, et al. Hypovitaminosis D-related myopathy in immigrant teenagers. *Neuropediatrics.* 2004. 35(5):290-292.
- 73. Block SR. Vitamin D deficiency is not associated with nonspecific musculoskeletal pain syndromes including fibromyalgia [letter]. *Mayo Clin Proc.* 2004. 79:1585-1591. Avail at: <http://www.mayoclinicproceedings.com/inside.asp?AID=2257&UID>. Access checked 3/12/08.
- 74. Al Faraj S, Al Mutairi K. Vitamin D deficiency and chronic low back pain in Saudi Arabia. *Spine* 2003. 28:177-179.
- 75. Huisman AM, White KP, Algra A, et al. Vitamin D levels in women with systematic lupus erythematosus and fibromyalgia. *J Rheumatol.* 2001. 28(11):2535-2539.
- 76. Glerup H, Mikkelsen K, Poulsen L, et al. Hypovitaminosis D myopathy without biochemical signs of osteomalacic bone involvement. *Calcif Tissue Int.* 2000b. 66:419-424.
- 77. Prabhala A, Garg R, Dandona P. Severe myopathy associated with vitamin D deficiency in

western New York. *Arch Intern Med.* 2000. 160(8):1199-1203.

- 78. McAlindon TE, Felson DT, Zhang Y, et al. Relation of dietary intake and serum levels of vitamin D to progression of osteoarthritis of the knee among participants in the Framingham Study. *Ann Int Med.* 1996. 125(5):353-359. Avail at: <http://www.annals.org/cgi/content/full/125/5/353>. Accessed 4/21/08
- 79. Gloth FM 3rd, Lindsay JM, Zelesnick LB, Greenough WB 3rd. Can vitamin D deficiency produce an unusual pain syndrome? *Arch Intern Med.* 1991. 151:1662-1664.
- 80. Boland R. Role of vitamin D in skeletal muscle function. *Endocr Rev.* 1986;7:434-438.
- 81. Haddad JG, Walgate J, Min C, Hahn TJ. Vitamin D metabolite-binding proteins in human tissue. *Biochem Biophys Acta.* 1976;444:921-925.
- 82. Schott G, Wills M. Muscle weakness in osteomalacia. *Lancet.* 1976;2:626-629.
- 83. Simpson RU, Thomas GA, Arnold AJ. Identification of 1,25-dihydroxyvitamin D3 receptors and activities in muscle. *J Biol Chem.* 1985. 260:8882-8891.
- 84. Pfeifer M, Begerow B, Minne HW, et al. Vitamin D status, trunk muscle strength, body sway, falls, and fractures among 237 postmenopausal women with osteoporosis. *Exp Clin Endocrinol Diabetes.* 2001. 109:87-92.
- 85. Bischoff HA, Stahelin HB, Dick W, et al. Effects of vitamin D and calcium supplementation on falls: a randomized controlled trial. *J Bone Miner Res.* 2003. 18:343-351.
- 86. Bischoff-Ferrari HA, Willett WC, Wong JB, et al. Fracture prevention with vitamin D supplementation: a meta-analysis of randomized controlled trials. *JAMA.* 2005. 293(18):2257-2264.
- 87. Bischoff-Ferrari HA, Giovannucci E, Willett WC, et al. Estimation of optimum serum concentrations of 25-hydroxyvitamin D for multiple health outcomes. *Am J Clin Nutr.* 2006. 84:18-28.
- 88. Dukas L, Bischoff HA, Lindpainter LS, et al. Alfacalcidol reduces the number of fallers in a community-dwelling elderly population with a minimum calcium intake of more than 500 mg daily. *J Am Geriatr Soc.* 2004. 52:230-236.
- 89. Sambrook PN, Chen JS, March LM, et al. Serum parathyroid hormone predicts time to fall independent of vitamin D status in a frail elderly population. *J Clin Endocrinol Metab.* 2004. 89:1572-1576.
- 90. Boxer RS, Dauser RA, Walsh SJ, et al. The association between vitamin D and inflammation with the 6-minute walk and frailty in patients with heart failure. *J Am Geriatr Soc.* 2008. 56:454-461.
- 91. Lee P, Chen R. Vitamin D as an analgesic for patients with type 2 diabetes and neuropathic pain. *Arch Intern Med.* 2008. 168(7):771-772.
- 92. D'Ambrosio D, Cippitelli M, Cocciolo MG, et al. Inhibition of IL-12 production by 1,25-Dihydroxyvitamin D3. *J Clin Invest.* 1998. 101:252-262.
- 93. Schleithoff SS, Zittermann A, Tenderich G, et al. Vitamin D supplementation improves cytokine profiles in patients with congestive heart failure: a double-blind randomized, placebo-controlled trial. *Am J Clin Nutr.* 2006. 83:754-759.
- 94. Van den Berghe, Van Roosbroeck D, Vanhove P, et al. Bone turnover in prolonged critical illness: effect of vitamin D. *J Clin Endocr Metab.* 2003. 88:4623-4632.
- 95. Thys-Jacobs S. Alleviation of migraine with therapeutic vitamin D and calcium. *Headache.* 1994a. 34(10):590-592.
- 96. Thys-Jacobs S. Vitamin D and calcium in menstrual migraine. *Headache.* 1994b;34(9):544-546.
- 97. Gloth FM 3rd, Alam W, Hollis B. Vitamin D vs broad spectrum phototherapy in the treatment of seasonal affective disorder. *J Nutr Health Aging.* 1999. 3(1):5-7.
- 98. Lansdowne AT, Provost SC. Vitamin D3 enhances mood in healthy subjects during winter. *Psychopharmacology.* 1998. 135(4):319-323.
- 99. Basaran S, Guzel R, Coskun-Benliday I, Guler-Uysal F. Vitamin D status: effects on quality of life in osteoporosis among Turkish women. *Qual Life Res.* 2007. 16(9):1491-1499.
- 100. Vieth R, Kimball S, Hu A, Walfish PG. Randomized comparison of the effects of the vitamin D3 adequate intake versus 100mcg (4000 IU) per day on biochemical responses and the wellbeing of patients. *Nutrition J.* 2004. 3:8. Avail at: <http://www.nutritionj>.

com/content/pdf/1475-2891-3-8.pdf. Accessed 5/14/08.

- 101. Smith GR, Collinson PO, Kiely PD. Diagnosing hypovitaminosis D: serum measurements of calcium, phosphate, and alkaline phosphatase are unreliable, even in the presence of secondary hyper-parathyroidism. *Rheumatol*. 2005. 32(4):684-689.
- 102. Nellen JFJB, Smulders YM, PH Jos Frissen, EH Slaats, J Silberbusch. Hypovitaminosis D in immigrant women: slow to be diagnosed. *BMJ*. 1996. 312:570-572.
- 103. Peach H, Compston JE, Vedi S, Horton LWL. Value of plasma calcium, phosphate, and alkaline phosphate measurements in the diagnosis of histological osteomalacia. *J Clin Pathol*. 1982;35:625-630. Avail at: <http://jcp.bmj.com/cgi/reprint/35/6/625>. Accessed 3/7/08.
- 104. Cooper MS, Gittoes NJL. Diagnosis and management of hypocalcemia. *BMJ*. 2008. 336:1298-1302.
- 105. Fitzpatrick L. Hypocalcemia: Diagnosis and Treatment – Chapter 7. 2002. Endotext.com. Avail at:
  - <http://www.endotext.org/parathyroid/parathyroid7/parathyroid7.htm>. Accessed 2/23/08.
- 106. Lyman D. Undiagnosed vitamin D deficiency in the hospitalized patient. *Am Fam Phys*. 2005. 71(2):299-304. Avail at: <http://www.aafp.org/afp/20050115/299.pdf>. Accessed 4/30/08.
- 107. Skugor M, Milas M. Hypocalcemia. 2004. Cleveland Clinic; Avail at: <http://www.clevelandclinicmeded.com/>. Accessed 2/23/08.
- 108. Peacey SR. Routine biochemistry in suspected vitamin D deficiency. *JR Soc Med*. 2004. 97:322-325. Avail at: <http://www.jrsm.org/cgi/reprint/97/7/322>. Accessed 3/8/08.
- 109. Zerwekh JE. Blood biomarkers of vitamin D status. *Am J Clin Nutr*. 2008. 87(suppl):1087S-1094S.
- 110. Singh RJ. Are clinical laboratories prepared for accurate testing of 25-hydroxy vitamin D? [letter]. *Clin Chem*. 2008. 54(1):221-223.
- 111. Masood H, Narang AP, Bhat IA, Shah GN. Persistent limb pain and raised serum alkaline phosphatase the earliest markers to subclinical hypovitaminosis D in Kashmir [abstract]. *Indian J Physiol Pharmacol*. 1989. 33(4):259-261.
- 112. Holick MF. Vitamin D deficiency as a contributor to multiple forms of chronic pain – reply II [letter]. *Mayo Clin Proc*. 2004a;79:694-709. Avail at: <http://www.mayoclinicproceedings.com/inside.asp?AID=2197&UID>. Accessed 3/7/08.
- 113. IOM (Institute of Medicine). Standing Committee on the Scientific Evaluation of Dietary Reference Intakes. Dietary Reference Intakes for Calcium, Phosphorus, Magnesium, Vitamin D, and Fluoride. Washington, DC: National Academy Press. 1997. 71-145.
- 114. DHHS (US Department of Health and Human Services and US Dept of Agriculture). Dietary Guidelines for Americans, 2005. 6th ed. Washington, DC: US Government Printing Office. 2005(Jan). Avail at: <http://www.health.gov/dietaryguidelines/dga2005/document/>. Accessed 5/18/08.
- 115. Armas LAG, Hollis BW, Heaney RP. Vitamin D2 is much less effective than vitamin D3 in humans. *J Clin Endocrinol Metab*. 2004. 89(11):5387-5391. Avail at: <http://jcem.endojournals.org/cgi/reprint/89/11/5387>. Accessed 4/23/08.
- 116. Barger-Lux MJ, Heaney RP, Dowell S, Chen TC, Holick MF. Vitamin D and its major metabolites: serum levels after graded oral dosing in healthy men. *Osteoporos Int*. 1998. 8(3):222-230.
- 117. Cranney A, Horsley T, O'Donnell S, et al. Effectiveness and safety of vitamin D in relation to bone health. *Evid Rep Technol Assess*. 2007. 158:1-235.
- 118. Trang HM, Cole DEC, Rubin LA, et al. Evidence that vitamin D3 increases serum 25-hydroxyvitamin D more efficiently than does vitamin D2. *Am J Clin Nutr*. 1998. 68:854-858. Avail at: <http://www.ajcn.org/cgi/reprint/68/4/854>. Accessed 4/23/08.
- 119. Vieth R, Bischoff-Ferrari H, Boucher BJ. The urgent need to recommend an intake of vitamin D that is effective. *Am J Clin Nutr*. 2007. 85:649-650.
- 120. Smith H, Anderson F, Raphael P, et al. Effect of annual intramuscular vitamin D on fracture risk in elderly men and women – a population-based, randomized, double-blind, placebo-controlled trial. *Rheumatology*. 2007. 46(12):1852-1857.
- 121. Hathcock JN, Shao A, Vieth R, Heaney R. Risk assessment for vitamin D. *Am J Clin Nutr*. 2007. 85:6-18.

- 122. Trivedi DP, Doll R, Khaw KT. Effect of four monthly oral vitamin D3 (cholecalciferol) supplementation on fractures and mortality in men and women living in the community: randomized double blind controlled trial. *Br Med J*. 2003. 326:469.
- 123. Romagnoli E, Mascia ML, Cipriani C, et al. Short and long term variations in serum calcitrophic hormones after a single very large dose of ergocalciferol (vitamin D2) or cholecalciferol (vitamin D3) in the elderly. *J Clin Endocrin Metab*. 2008. online publication ahead of print; doi:10.1210/jc.2008-0350.
- 124. Hollis BW, MacFarlane G, Valcour A. Clinical significance and assessment of 25 OH vitamin D. Burlington, NC: LabCorp; 2005. Avail at: [http://www.labcorp.com/cme/pdf/Vitamin\\_D\\_Journal\\_Article.pdf](http://www.labcorp.com/cme/pdf/Vitamin_D_Journal_Article.pdf). Accessed 4/16/08.
- 125. Barrueto F Jr, Wang-Flores HH, Howland MA, Hoffman RS, Nelson LS. Acute vitamin D intoxication in a child. *Pediatrics*. 2005. 116:e453-e456.
- 126. Blank S, Scanlon KS, Sinks TH, Lett S, Falk H. An outbreak of hypervitaminosis D associated with the over fortification of milk from a home-delivery dairy. *Am J Pub Health*. 1995. 85(5):656-659.
- 127. Carroll MF, Schade DS. A practical approach to hypercalcemia. *Am Fam Physician*. 2003. 67:1959-1966. Avail at: <http://www.aafp.org/afp/20030501/1959.html>. Accessed 4/21/08.
- 128. Klontz KC, Acheson DW. Dietary supplement-induced vitamin D intoxication [letter]. *New Eng J Med*. 2007. 357:308-309.
- 129. Koutkia P, Chen TC, Holick MF. Vitamin D intoxication associated with an over-the-counter supplement. *New Engl J Med*. 2001. 345(1):65-66.
- 130. Todd MA, Bailey RR, Espiner EA, Lynn KL. Vitamin D2 for the treatment of chilblains – a cautionary tale. *N Z Med J*. 1987;100:465.
- 131. Vieth R, Pinto TR, Reen BS, Wong MM. Vitamin D poisoning by table sugar. *Lancet*. 2002. 359:672.
- 132. Jacobus CH, Holick MF, Shao Q, et al. Hypervitaminosis D associated with drinking milk. *New Engl J Med*. 1992;326:1173-1177.
- 133. ODS (Office of Dietary Supplements; US National Institutes of Health). Dietary Supplement Fact Sheet: Calcium. Updated Sept 23, 2005. Avail at: [http://dietary-supplements.info.nih.gov/factsheets/Calcium\\_pf.asp](http://dietary-supplements.info.nih.gov/factsheets/Calcium_pf.asp). Accessed 4/16/08
- 134. Adams JS, Lee G. Gains in bone mineral density with resolution of vitamin D intoxication. *Ann Int Med*. 1997. 127(3):203-206.
- 135. Pettifor JM, Bikle DD, Cavaleros M. Serum levels of free 1,25-dihydroxyvitamin D in vitamin D toxicity. *Ann Int Med*. 1995. 122(7):511-513.
- 136. Bronstein AC, Spyker DA, Cantilene Jr LR, et al. 2006 Annual Report of the American Association of Poison Control Centers National Poison Data System (NPDS). *Clinical Toxicology*. 2007. 45(8):815-917. Avail at: <http://www.aapcc.org/poison1.htm>. Accessed 2/22/08.
- 137. Bringhurst RF, Bemay MB, Krune SM, Kronenberg HM. Bone and mineral metabolism in health and disease. In: Kasper DL, et al. eds. *Harrison's Principles of Internal Medicine*. 16th Ed. New York: McGraw-Hill; 2005: 2238-2249.

First published on: July 1, 2008

**Source URL:** <http://www.practicalpainmanagement.com/treatments/nutraceutical/vitamin-d-chronic-pain>

**Links:**

[1] [http://www.practicalpainmanagement.com/sites/default/files/imagecache/lightbox-large/import\\_files/html/P0807F02/fig1.png](http://www.practicalpainmanagement.com/sites/default/files/imagecache/lightbox-large/import_files/html/P0807F02/fig1.png)

[2] [http://www.practicalpainmanagement.com/sites/default/files/imagecache/lightbox-large/import\\_files/html/P0807F02/table3.jpg](http://www.practicalpainmanagement.com/sites/default/files/imagecache/lightbox-large/import_files/html/P0807F02/table3.jpg)

[3] [http://www.practicalpainmanagement.com/sites/default/files/imagecache/lightbox-large/import\\_files/html/P0807F02/table3\\_02.jpg](http://www.practicalpainmanagement.com/sites/default/files/imagecache/lightbox-large/import_files/html/P0807F02/table3_02.jpg)

[4] #fieldset