

Medication-overuse headache: epidemiology, diagnosis and treatment

Espen Saxhaug Kristoffersen and Christofer Lundqvist

Ther Adv Drug Saf

2014, Vol. 5(2) 87–99

DOI: 10.1177/
2042098614522683

© The Author(s), 2014.
Reprints and permissions:
[http://www.sagepub.co.uk/
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

Abstract: Medication-overuse headache (MOH) is one of the most common chronic headache disorders and a public health problem with a worldwide prevalence of 1–2%. It is a condition characterized by chronic headache and overuse of different headache medications, and withdrawal of the overused medication is recognised as the treatment of choice. However, the strategy for achieving withdrawal is, at present, based on expert opinion rather than scientific evidence, partly due to the lack of randomised controlled studies. This narrative review investigates different aspects of epidemiology, diagnosis, risk factors and pathogenesis as well as management for MOH. We suggest that the first step in the treatment of MOH should be carried out in general practice and should focus primarily on detoxification. For most patients, both prevention and follow up after detoxification can also be performed in general practice, thus freeing resources for referral of more complicated cases to headache clinics and neurologists. These suffering patients have much to gain by an earlier treatment-focused approach lower down on the treatment ladder.

Keywords: chronic headache, dependence, drug safety, medication overuse, migraine

Introduction

Headache is very common and usually occurs episodically, but 3–4% of the general population have chronic headache, defined as 15 or more days of headache per month [Castillo *et al.* 1998; Lantéri-Minet *et al.* 2003; Grande *et al.* 2008; Aaseth *et al.* 2008; Headache Classification Committee of the International Headache Society, 2013]. Headache disorders as a whole are a major public health concern given the large amount of associated disability and financial costs both to the individual and to society [Stovner *et al.* 2007; Jensen and Stovner, 2008; Linde *et al.* 2012; Vos *et al.* 2012]. Patients with chronic headache represent a large population within primary care and in neurologist settings [Patterson and Esmonde, 1993; Latinovic *et al.* 2006; Ridsdale *et al.* 2007]. Headache is often treated with analgesics and is the most common reason for analgesic use in the general population [Eggen, 1993; Antonov and Isacson, 1998; Zwart *et al.* 2003, 2004; Mehuys *et al.* 2012]. In Scandinavia and Scotland, 20–40% had used analgesics over the last 14 days and, in 2007, sales figures of over-the-counter (OTC) analgesics in Denmark were

extrapolated to correspond to almost 8% of the general Danish population taking the highest recommended daily dose every day for a whole year [Eggen, 1993; Antonov and Isacson, 1998; Porteous *et al.* 2005; Hargreave *et al.* 2010].

Inappropriate use of symptomatic medication for headaches may paradoxically lead to medication-overuse headache (MOH) [Headache Classification Committee of the International Headache Society, 2013]. This is a condition characterized by chronic headache and overuse of different acute headache medications. Withdrawal headache after excessive intake of ergotamine was described in the early 1950s and 1960s, and from the 1980s studies have shown that frequent intake of all symptomatic headache medication may transform episodic headache to frequent headache [Peters and Horton, 1950, 1951; Horton and Peters, 1963; Kudrow, 1982; Diener *et al.* 1989; Mathew *et al.* 1990]. MOH patients, like other chronic headache sufferers, experience reduced quality of life compared with those do not suffer from headache and episodic headache [Colás *et al.* 2004; Lantéri-Minet *et al.* 2011].

Correspondence to:
Espen Saxhaug Kristoffersen, MD
Department of General Practice, Institute of Health and Society, University of Oslo, Oslo, and Research Centre, Akershus University Hospital, Lørenskog, Norway
e.s.kristoffersen@medisin.uio.no
Christofer Lundqvist, MD, PhD
Research Centre, Akershus University Hospital, Lørenskog, Norway; Institute of Clinical Medicine, Campus Akershus University Hospital, University of Oslo, Nordbyhagen, Norway, and Department of Neurology, Akershus University Hospital, Nordbyhagen, Norway

The treatment of MOH is often complex and withdrawal of the overused medication is recognised as the treatment of choice [Evers and Jensen, 2011]. Though several reviews on MOH have been published previously [Diener and Limmroth, 2004; Katsarava and Jensen, 2007; Rossi *et al.* 2009; Evers and Marziniak, 2010; Russell and Lundqvist, 2012; Katsarava and Obermann, 2013], this review will attempt to give an up-to-date focus on epidemiology, diagnosis and different aspects of the management, treatment and prevention of MOH.

Classification of medication-overuse headache

The International Classification of Headache Disorders 3rd beta edition (ICHD-III β) divides headaches into primary and secondary forms [Headache Classification Committee of the International Headache Society, 2013]. Primary headaches are idiopathic disorders without other known causes, whereas secondary headaches are headaches assumed to be caused by other illnesses or external factors (e.g. trauma, surgery, toxic effects of substances or medications or infection). According to the ICHD-III β :

MOH is headache occurring on 15 or more days per month developing as a consequence of regular overuse of acute or symptomatic headache medication (on 10 or more, or 15 or more days per month, depending on the medication) for more than 3 months. It usually, but not invariably, resolves after the overuse is stopped (Box 1).

MOH is classified as a secondary chronic headache, but whether MOH is a primary or secondary headache is still under debate, and the concept of medication overuse in other secondary headaches is unclear [Sun-Edelstein *et al.* 2009; Grazzi and Bussone, 2012].

Epidemiological and socioeconomic aspects of MOH

The prevalence of MOH in the general population is 1–2% [Lu *et al.* 2001; Colás *et al.* 2004; Zwart *et al.* 2004; Wiendels *et al.* 2006; Grande *et al.* 2008; Aaseth *et al.* 2008; Straube *et al.* 2010; Jonsson *et al.* 2011]. The male : female ratio is 1 : 3–4 and it is most prevalent in the forties [Straube *et al.* 2010; Jonsson *et al.* 2011]. The prevalence decreases with older age and, among people over 65 years in Taiwan, the prevalence was 1.0%

Box 1. International Classification of Headache Disorders, 3rd beta edition (ICHD-III β) criteria for medication-overuse headache [Headache Classification Committee of the International Headache Society, 2013].

- A. Headache present on >15 days/month.
- B. Regular overuse for >3 months of one or more drugs that can be taken for acute and/or symptomatic treatment of headache.
- C. Headache has developed or markedly worsened during medication overuse.

(For simple analgesics and for combination of acute medications, the intake must be 15 days or more per month for triptans, ergotamins, opioids and combination analgesics; 10 days per month is enough to get the diagnosis of MOH.)

[Wang *et al.* 2000]. The prevalence of MOH in children and adolescents has been suggested to be 0.3–0.5% [Dyb *et al.* 2006; Wang *et al.* 2006]. MOH generally starts earlier in life than other types of chronic headache [Colás *et al.* 2004].

While OTC drugs are the most commonly overused headache medications in primary care, secondary and tertiary care have a greater proportion of MOH patients who overuse more potent, centrally acting drugs [Lu *et al.* 2001; Meskunas *et al.* 2006; Zeeberg *et al.* 2006; Scher *et al.* 2010; Jonsson *et al.* 2011, Kristoffersen *et al.* 2012; 2013]. The drugs involved in MOH change over time and from region to region [Meskunas *et al.* 2006]. In the US it has been found that 23% of chronic headache sufferers used acute medication on a daily basis in contrast to only 9% in the Norwegian general population [Scher *et al.* 2010; Kristoffersen *et al.* 2012]. In contrast to previously, ergotamine is no longer a large problem in western Europe. However, triptans are now among the most common causes of MOH in the western world. It has been suggested that the mean critical duration of overuse is shortest for triptans (1.7 years), longer for ergotamine (2.7 years) and longest for simple analgesics (4.8 years). The mean critical monthly intake frequency in the same study was lowest for triptans, higher for ergotamine and highest for analgesics [Limmroth *et al.* 2002].

Risk factors

Many psychosocial and socioeconomic factors are associated with MOH. However, it is hard to ascertain if these are directly or indirectly associated as these findings are mainly based on

cross-sectional studies. Thus, many of these factors may merely be markers of a complex situation since many aspects of life may be affected by having chronic headache, as with other chronic conditions.

As for other frequent headaches, MOH patients tend to have a low socioeconomic status with low income and education, but it is uncertain whether this may be a cause of or an effect of headache [Hagen *et al.* 2002; Atasoy *et al.* 2005; Wiendels *et al.* 2006; Jonsson *et al.* 2011, 2012]. A high prevalence of smoking, high body mass index and sleeping problems have also been found among MOH patients [Wiendels *et al.* 2006; Straube *et al.* 2010]. Depression and anxiety was more common among MOH patients than among people with episodic migraine, but in another study this was related to the headache frequency rather than headache diagnosis [Zwart *et al.* 2003; Radat *et al.* 2005]. The risk of developing MOH is greater in individuals with a family history of MOH or other substance abuse [Cevoli *et al.* 2009].

Data from a population-based longitudinal study suggested that those who used analgesics daily or weekly at baseline had a higher risk of developing chronic headache 11 years later [Zwart *et al.* 2003]. This would seem to support a causative role of medication overuse in generating MOH.

A more recent study identified several risk factors for MOH among people with chronic headache (11 years follow up) [Hagen *et al.* 2012]. Regular use of tranquilizers, combination of chronic musculoskeletal and gastrointestinal complaints, and increased Hospital Anxiety and Depression Scale, as well as smoking and physical inactivity increased the risk for MOH. The study was extensive and included over 25,000 people at risk for chronic headache and MOH. However, the mentioned risk factors were just found in a minority of all the MOH patients, and may thus reflect the complex situation for specific subgroups of MOH patients rather than for the MOH patient in general.

Previous primary headache such as migraine and tension-type headache are also risk factors and seems to be required for the development of MOH [Bahra *et al.* 2003; Colás *et al.* 2004; Bigal and Lipton, 2008].

Societal consequences of MOH

MOH is probably the most the most costly headache disorder for both society and the sufferer

[Linde *et al.* 2012]. In a recent assessment of direct and indirect costs of headache disorders in Europe, indirect loss due to reduced productivity and absenteeism accounts for about 90% of the costs [Linde *et al.* 2012]. The individual costs of MOH were higher than for migraine, and the total national costs for headache disorders in some countries were estimated to be higher for MOH than for migraine. Total national costs for MOH were estimated at €5–10 billion in Italy, Spain and France [Linde *et al.* 2012]. Thus, the worldwide personal and economic costs are enormous.

Pathogenesis

Approximately half of those with headache on more than 15 days per month have MOH [Grande *et al.* 2008; Aaseth *et al.* 2008]. Most headache experts regard the association between overuse of acute medication and development of MOH as causal [Bigal and Lipton, 2009; Evers and Jensen, 2011]. Improvement for two-thirds to three-quarters of patients upon removal of the overused medication supports its causative role in generating or maintaining a chronic headache.

However, it is still a matter of debate whether the overuse is a consequence of living with chronic headache or the other way round [Dodick, 2002; Tepper, 2002]. Furthermore, not all headache patients with medication overuse develop MOH, and the mechanism how chronic exposure to abortive medication leads to MOH remains unclear.

Virtually all acute headache medication may cause MOH and since the different medications have different pharmacological actions, it is unlikely that MOH is caused by the specific action of any single agent. Mechanisms may differ from one class of overused drug to another and different possible pathogeneses have been suggested. It is of course possible that there is a common, but still unknown, mechanism by which pharmacologically different medications leads to MOH. However, at present it is possible only to describe mechanisms (mostly from preclinical studies) that appear to be associated with, or may predispose people to develop MOH.

A pre-existent headache disorder seems to be required to develop MOH [Bahra *et al.* 2003]. Migraine and tension-type headache have a higher potential for developing MOH than other

primary headaches, but also patients with cluster headaches may develop MOH [Colás *et al.* 2004; Paemeleire *et al.* 2006; Bigal and Lipton, 2008]. However, MOH does not develop in persons without a history of headache when medication is taken regularly for other conditions such as arthritis or inflammatory bowel disease [Wilkinson *et al.* 2001; Bahra *et al.* 2003]. Thus, a connection between headache-specific pain pathways and headache medication effects seems to be a central factor in generating a more chronic pain.

A hereditary susceptibility to MOH has been suggested as the risk of developing MOH is greater in individuals with a family history of MOH or other substance abuse [Cevoli *et al.* 2009]. A few small-scale studies have found some molecular genetic factors that are possibly associated with MOH, but these results are from small studies in selected groups and the generalizability of the findings is difficult to ascertain [Cevoli *et al.* 2006; Di Lorenzo *et al.* 2007, 2009]. Thus, further studies are required.

Alteration of cortical neuronal excitability, central sensitization involving the trigeminal nociceptive system, and changes in serotonergic and dopaminergic expressions and pathways including the endocannabinoid system have been suggested to play a part in the pathophysiology of MOH [Cevoli *et al.* 2006; Di Lorenzo *et al.* 2009; Cupini *et al.* 2010; De Felice *et al.* 2010; Meng *et al.* 2011; Bongsebandhu-Phubhakdi and Srikiatkachorn, 2012].

Low serotonin (5-HT) levels with reduction of 5-HT in platelets and upregulation of a pronociceptive 5-HT_{2A} receptor have been demonstrated in MOH [Srikiatkachorn *et al.* 1998; Srikiatkachorn and Anthony, 2006]. A higher frequency of cortical spreading depression (CSD) has been found in animals with low 5-HT levels suggesting an association with sensitization processes [Supornsilpchai *et al.* 2006; Le Grand *et al.* 2011]. Furthermore, chronic, but not acute, paracetamol administration led to an increase in CSD frequency in another rat model which may indicate that chronic analgesic exposure leads to hyperexcitability in cortical neurons and an increase in CSD [Supornsilpchai *et al.* 2010]. A 5-HT_{2A} receptor antagonist blocked this increased CSD susceptibility in the rats which had been exposed to chronic paracetamol [Supornsilpchai *et al.* 2010].

Chronic use of opioids and triptans has been shown to increase calcitonin gene related peptide (CGRP) levels which is involved in neurogenic inflammation and headache pain [Belanger *et al.* 2002; De Felice *et al.* 2010]. In addition, chronic morphine infusion may alter the diffuse noxious inhibitory controls (DNICs) and impaired DNICs are also found in MOH [Perrotta *et al.* 2010].

Another important phenomenon related to MOH is central sensitization which has for a long time been suggested to play an important role and has recently been described clinically in MOH patients with normalization after withdrawal of the overused medication [Ayzenberg *et al.* 2006; Munksgaard *et al.* 2013].

The endocannabinoid system is involved in modulating pain and plays a role in the common neurobiological system underlying drug addiction and reward [Cupini *et al.* 2008]. Both an endocannabinoid membrane transporter and levels of endocannabinoids in platelets were reduced in MOH and chronic migraine patients compared to controls [Cupini *et al.* 2008; Rossi *et al.* 2008].

In MOH patients, increased levels of orexin-A and corticotrophin-releasing hormone were found in the cerebrospinal fluid compared with patients with chronic migraine and these levels were correlated to monthly drug intake [Sarchielli *et al.* 2008]. In addition, neuroimaging studies suggest changes in the orbitofrontal cortex and the mesocorticolimbic dopamine circuit [Fumal *et al.* 2006; Ferraro *et al.* 2012].

In conclusion, the complex pathophysiology behind MOH is still only partly known. However, it is clear that many of these phenomena are similar to and thus may involve mechanisms seen in dependence processes [Calabresi and Cupini, 2005; Cupini *et al.* 2010] and it is equally clear that more research in these areas is needed.

Medication dependence or not?

Although triptans and simple analgesics are not regarded as psychotropic agents, patients with MOH do share some characteristics with dependence and some authors have advocated the division of MOH into two subgroups depending on the type of overused medication and on co-morbidity [Lake, 2006; Saper and Lake, 2006].

Codeine and opioids are not recommended in the treatment of tension-type headache or migraine. Regardless of this, it is well known that many MOH patients use these agents [Evers *et al.* 2009; Bendtsen *et al.* 2010]. Furthermore, many countries have codeine (which is metabolised to opioids) and caffeine-containing analgesics available as OTC drugs. Codeine, opioids and caffeine are known to be psychotropic drugs; thus abuse and dependence on headache drugs may be a problem [Abbott and Fraser, 1998; Cooper, 2013a, 2013b].

Theoretical considerations also show many similarities between MOH and drug addiction [Calabresi and Cupini, 2005]. Previous studies from Norway have revealed that MOH can easily be detected in a population using a screening instrument for behavioural dependence – the Severity of dependence scale (SDS) [Gossop *et al.* 1995; Grande *et al.* 2009; Lundqvist *et al.* 2010]. The SDS has high sensitivity, specificity, positive and negative predictive values for detecting persons with MOH among chronic headache patients [Grande *et al.* 2009; Lundqvist *et al.* 2010, 2011]. In addition, the SDS score has been shown to predict likelihood of successful detoxification in a general population [Lundqvist *et al.* 2012]. The SDS has not been validated against other measurements of dependency in MOH sufferers.

In two studies, approximately 70% of MOH patients fulfilled criteria for dependence according to the Diagnostic and Statistical Manual of Mental Disorders IV (DSM-IV) [Fuh *et al.* 2005; Radat *et al.* 2008]. However, the use of DSM-IV criteria in patients with MOH has been criticised since it may overestimate dependence [Lauwerier *et al.* 2011]. Another study found that the dependency score based on the Leeds Dependency Questionnaire (LDQ) was similarly increased in MOH patients and illegal drug addicts [Ferrari *et al.* 2006]. However, in contrast to drug addicts, it was not the type of drug that was most important for the MOH patients but the effect of the drug [Ferrari *et al.* 2006].

Based on today's knowledge, it is impossible to ascertain whether dependency-like behaviour seen in those with MOH represents a real dependence or whether it is a form of 'pseudo-addiction' secondary to frequent headache [Saper *et al.* 2005; Lake, 2006; Saper and Lake, 2006; Radat and Lantéri-Minet, 2010; Fuh and Wang, 2012].

Treatment

MOH is a heterogeneous and complex condition which includes both simple and complicated overuse [Saper *et al.* 2005; Lake, 2006; Rossi *et al.* 2011]. A further challenge in the treatment is the fact that there is no worldwide consensus for the management of these patients other than that termination of medication overuse is desirable. How this should be addressed is frequently discussed. This may be mainly due to the lack of controlled studies, but also reflects the differences in use of the headache classification and nondocumented treatment strategies across the world. Although debated, withdrawal of the overused medication(s) is the treatment of choice, since withdrawal of the overused medication(s) in most cases leads to an improvement of the headache [Linton-Dahlöf *et al.* 2000; Zeeberg *et al.* 2006; Evers and Jensen, 2011; Diener, 2012; Olesen, 2012].

Most patients experience withdrawal symptoms lasting 2–10 days after detoxification. The most common symptom is an initial worsening of the headache, accompanied by various degrees of nausea, vomiting, hypotension, tachycardia, sleep disturbances, restlessness, anxiety and nervousness. The duration of withdrawal headaches have been found to vary with different drugs, being shorter in patients overusing triptans (~4 days) than in ergotamine (~7 days) or analgesics (~10 days) [Katsarava *et al.* 2001].

The procedures for detoxification vary substantially and include both in-patient (2 days to 2 weeks) and out-patient withdrawal [Krymchantowski and Moreira, 2003; Grazzi *et al.* 2008; Creac'h *et al.* 2011; Gaul *et al.* 2011; Rossi *et al.* 2011, Munksgaard *et al.* 2012]. The different strategies include: just simple advice; multidisciplinary approaches; use of antiemetics, tranquilizers, neuroleptics, rescue medication (another analgesic than the overused); intravenous hydration; and/or administration of oral, nasal or intravenous ergotamines. Steroids have for a long time been expected to alleviate withdrawal headache in the acute phase, but two placebo-controlled studies did not find prednisolone (60 or 100 mg for 5 days) superior to placebo [Boe *et al.* 2007; Rabe *et al.* 2013]. However, it may be useful in subgroups.

Regardless of the strategy, the main aims of the treatment are:

- (i) Withdrawal of the overused drug(s)
- (ii) To provide the patient with pharmacological and nonpharmacological support
- (iii) To prevent relapse.

Clinical studies of treatment results from headache centres often have treatment success rates of around 70%. These results are commonly based on in-patient treatment, rescue medication and continued support. In addition, the 70% success rate is based on very variable outcome measures and therefore difficult to compare [Hagen *et al.* 2010]. No randomised controlled trials have been conducted in the general population or in primary care. However, some studies have suggested effect of simple advice for MOH in both a research setting in the general population and in a headache clinic [Rossi *et al.* 2006, 2011, 2013; Grande *et al.* 2011]. In a previous population-based study from Norway, simple information on medication use led to improvement with 42% of patients reverting to episodic headache and 76% being free of medication overuse after 1.5 years. The study was observational and lacked a control group, but the effect of simple information was supported by the fact that the participants had had MOH for 8–18 years prior to the interview during which the information was given [Grande *et al.* 2011]. Two Italian studies from neurologist settings have reported 78–92% of simple MOH patients and 60% of complicated MOH to be without chronic headache and medication overuse two months after simple advice [Rossi *et al.* 2006, 2011, 2013].

There is still some debate as to whether or not initially to detoxify MOH patients and whether prophylactic medication should be initiated immediately at withdrawal or after completed withdrawal therapy [Evers and Jensen, 2011; Diener, 2012; Olesen, 2012].

Placebo-controlled studies of prophylactic medication (onabotulinumtoxin A and topiramate) for chronic migraine with medication overuse have found a significant reduction in migraine and headache days per month compared with placebo [Diener *et al.* 2007, 2009, 2010; Dodick *et al.*, 2010; Aurora *et al.* 2010; Sandrini *et al.* 2011]. However, these results are not superior to detoxification without prophylactic medication in both simple and complicated MOH in the general population and neurologist setting, or in treatment-resistant MOH patients in a tertiary headache centre [Grande *et al.* 2011; Rossi *et al.* 2011; Munksgaard *et al.* 2012].

Therefore, based on today's knowledge we suggest that initial withdrawal is the treatment of choice. In addition, prophylactic headache medication should be restricted to patients that do not benefit sufficiently from simple advice or other means of withdrawal of medication overuse for the following reasons:

1. Removing offending medication is a logical primary aim for patients with headache partly due to medication side effects.
2. Removal of the distorting effect of medication overuse and assessment in an overuse-free state may be expected to show the characteristics of remaining headache and enable a rational choice of prophylactics, should they be needed.
3. A period free of medication overuse has been suggested to lead to recovery of prophylactic responsiveness [Zeeberg *et al.* 2006].

In analogy with other substance overuse, it is an aim in itself to achieve a behavioural change from 'have pain – take tablet' thinking to other, alternative and in the long run less detrimental ways of handling headache. Based on our clinical experience most patients overusing simple analgesics as well as codeine-containing combination medications and triptans manage abrupt detoxification from their offending medication without tapering. Patients overusing heavier drugs with physical abstinence profiles may be a different matter. However, in Norway as in most countries in Europe, the prior drug classes clearly dominate [Colás *et al.* 2004; Zeeberg *et al.* 2006; Jonsson *et al.* 2011; Kristoffersen *et al.* 2012, 2013].

Based on the lacking evidence base regarding detoxification strategies we are presently completing a randomised controlled trial of a behavioural-based, nonpharmacological short intervention in primary care. The results of this trial may contribute more firm evidence regarding whether detoxification in primary care can be achieved in primary out-patient care and whether it leads to lasting improvement of the headache situation of these patients [Kristoffersen *et al.* 2012].

Follow up and relapse

Studies from clinical settings have reported a 20–40% relapse rate of detoxified patients within the first year after withdrawal. Only few relapse after 12 months [Katsarava *et al.* 2003, 2005;

Grazzi *et al.* 2004; Zidverc-Trajkovic *et al.* 2007; Rossi *et al.* 2008; Boe *et al.* 2009a, 2009b; Andrasik *et al.* 2010]. There are conflicting results regarding at what time during the first year patients relapse with some suggesting that most patients relapse within the first 6 months while others suggest between 6 and 12 months [Katsarava *et al.* 2003, 2005; Grazzi *et al.* 2004; Ghiotto *et al.* 2009; Andrasik *et al.* 2010]. The literature is not clear regarding to what degree pre-existent headache type or type of overused medication predicts successful withdrawal [Limmroth *et al.* 2002; Katsarava *et al.* 2003, 2005; Grazzi *et al.* 2004; Zidverc-Trajkovic *et al.* 2007; Rossi *et al.* 2008; Boe *et al.* 2009a, 2009b; Andrasik *et al.* 2010].

In addition, these results on relapse should be compared with some caution, since the studies varied in the use of different headache classification systems, different withdrawal and prophylaxis as well as follow up and criteria for improvement.

Prevention

Some studies have found that most MOH patients do not know about the relationship between medication overuse and headache chronification [Bekkelund and Salvesen, 2002; Rossi *et al.* 2006; Jonsson *et al.* 2012]. The authors suggest that it is possible that the patients had been informed, but that they did not remember or had not fully understood the information. A brochure on medication overuse was preventive of development of MOH in people with migraine and frequent medication use [Fritsche *et al.* 2010].

Since most MOH patients have been in contact with their general practitioner (GP) for headache (80%), and almost half have had such contact in the previous year, primary care is probably the ideal setting for prevention and treatment of MOH [Jonsson *et al.* 2012; Kristoffersen *et al.* 2012, 2013]. Therefore, the GP has a key role in providing patient education and, in some cases, prophylactic headache medication before headaches become chronic.

It would be worthwhile to consider whether these MOH patients may benefit as much from a GP as they do from a neurologist. Taking care of uncomplicated cases in primary care may also free more resources for referrals to neurologists for complicated cases. Most follow-up studies are conducted in tertiary care centres and may reflect a selected

population. However, one Norwegian study supports that patients initially detoxified as inpatients, can be followed up by their GP [Boe *et al.* 2009].

The improvement after withdrawal of the overused medication may potentially be augmented by a greater awareness by doctors, pharmacists and society in general regarding the dangers associated with inappropriate use of painkillers for headache. Considering that possibly anyone with primary episodic headache may be at risk of developing MOH, the number of people at risk is high. New information campaigns and strategies to target the people under risk have to be developed since the potential benefit of information and prevention of MOH is thus also high.

Conclusion

MOH is a worldwide public health problem. The treatment may be complex, but improvements seen in two out of three MOH patients after withdrawal suggest detoxification to be the treatment of first choice. Prevention as well as primary detoxification is possible and should probably be attempted in primary care. The gain from treating patients with MOH is potentially high, and may lead to substantial economic savings for society as well as large benefits for the individual patients.

Funding

The authors received research funding from the University of Oslo, the Research Centre at Akershus University Hospital and the South-Eastern Norway Regional Health Authority.

Conflict of interest statement

The authors declare no conflicts of interest in preparing this article.

References

- Aaseth, K., Grande, R., Kvaerner, K., Gulbrandsen, P., Lundqvist, C. and Russell, M. (2008) Prevalence of secondary chronic headaches in a population-based sample of 30-44-year-old persons. The Akershus Study of Chronic Headache. *Cephalalgia* 28: 705-713.
- Abbott, F. and Fraser, M. (1998) Use and abuse of over-the-counter analgesic agents. *J Psychiatry Neurosci* 23: 13-34.
- Andrasik, F., Grazzi, L., Usai, S., Kass, S. and Bussone, G. (2010) Disability in chronic migraine with medication overuse: treatment effects through 5 years. *Cephalalgia* 30: 610-614.

- Antonov, K. and Isacson, D. (1998) Prescription and nonprescription analgesic use in Sweden. *Ann Pharmacother* 32: 485–494.
- Atasoy, H., Unal, A., Atasoy, N., Emre, U. and Sumer, M. (2005) low income and education levels may cause medication overuse and chronicity in migraine patients. *Headache* 45: 25–31.
- Aurora, S., Dodick, D., Turkel, C., Degryse, R., Silberstein, S., Lipton, R. *et al.* (2010) OnabotulinumtoxinA for treatment of chronic migraine: results from the double-blind, randomized, placebo-controlled phase of the Preempt 1 trial. *Cephalalgia* 30: 793–803.
- Ayzenberg, I., Obermann, M., Nyhuis, P., Gastpar, M., Limmroth, V., Diener, H. *et al.* (2006) Central sensitization of the trigeminal and somatic nociceptive systems in medication overuse headache mainly involves cerebral supraspinal structures. *Cephalalgia* 26: 1106–1114.
- Bahra, A., Walsh, M., Menon, S. and Goadsby, P. (2003) Does chronic daily headache arise de novo in association with regular use of analgesics? *Headache* 43: 179–190.
- Bekkelund, S. and Salvesen, R. (2002) Drug-associated headache is unrecognized in patients treated at a neurological centre. *Acta Neurol Scand* 105: 120–123.
- Belanger, S., Ma, W., Chabot, J. and Quirion, R. (2002) expression of calcitonin gene-related peptide, substance p and protein kinase c in cultured dorsal root ganglion neurons following chronic exposure to mu, delta and kappa opiates. *Neuroscience* 115: 441–453.
- Bendtsen, L., Evers, S., Linde, M., Mitsikostas, D., Sandrini, G., Schoenen, J. *et al.* (2010) EFNS guideline on the treatment of tension-type headache – report of an EFNS task force. *Eur J Neurol* 17: 1318–1325.
- Bigal, M. and Lipton, R. (2008) excessive acute migraine medication use and migraine progression. *Neurology* 71: 1821–1828.
- Bigal, M. and Lipton, R. (2009) overuse of acute migraine medications and migraine chronification. *Curr Pain Headache Rep* 13: 301–307.
- Boe, M., Mygland, A. and Salvesen, R. (2007) prednisolone does not reduce withdrawal headache. *Neurology* 69: 26–31.
- Boe, M., Salvesen, R. and Mygland, A. (2009a) Chronic daily headache with medication overuse: a randomized follow-up by neurologist or PCP. *Cephalalgia* 29: 855–863.
- Boe, M., Salvesen, R. and Mygland, A. (2009b) Chronic daily headache with medication overuse: predictors of outcome 1 year after withdrawal therapy. *Eur J Neurol* 16: 705–712.
- Bongsebandhu-Phubhakdi, S. and Srikiatkachorn, A. (2012) Pathophysiology of medication-overuse headache: implications from animal studies. *Curr Pain Headache Rep* 16: 110–115.
- Calabresi, P. and Cupini, L. (2005) Medication-overuse headache: similarities with drug addiction. *Trends Pharmacol Sci* 26: 62–68.
- Castillo, J., Munoz, P., Guitera, V. and Pascual, J. (1998) Kaplan Award 1998. Epidemiology of chronic daily headache in the general population. *Headache* 39: 190–196.
- Cevoli, S., Mochi, M., Scapoli, C., Marzocchi, N., Pierangeli, G., Pini, L. *et al.* (2006) A genetic association study of dopamine metabolism-related genes and chronic headache with drug abuse. *Eur J Neurol* 13: 1009–1013.
- Cevoli, S., Sancisi, E., Grimaldi, D., Pierangeli, G., Zanigni, S., Nicodemo, M. *et al.* (2009) Family history for chronic headache and drug overuse as a risk factor for headache chronification. *Headache* 49: 412–418.
- Colás, R., Muñoz, P., Temprano, R., Gómez, C. and Pascual, J. (2004) Chronic daily headache with analgesic overuse: epidemiology and impact on quality of life. *Neurology* 62: 1338–1342.
- Cooper, R. (2013a) ‘I can’t be an addict. I am.’ Over-the-counter medicine abuse: a qualitative study. *BMJ Open* 3:e002913
- Cooper, R. (2013b) Over-the-counter medicine abuse – a review of the literature. *J Subst Use* 18: 82–107.
- Creac’h, C., Frappe, P., Cancade, M., Laurent, B., Peyron, R., Demarquay, G. *et al.* (2011) In-patient versus out-patient withdrawal programmes for medication overuse headache: a 2-year randomized trial. *Cephalalgia* 31: 1189–1198.
- Cupini, L., Costa, C., Sarchielli, P., Bari, M., Battista, N., Eusebi, P. *et al.* (2008) Degradation of endocannabinoids in chronic migraine and medication overuse headache. *Neurobiol Dis* 30: 186–189.
- Cupini, L., Sarchielli, P. and Calabresi, P. (2010) Medication overuse headache: neurobiological, behavioural and therapeutic aspects. *Pain* 150: 222–224.
- De Felice, M., Ossipov, M., Wang, R., Lai, J., Chichorro, J., Meng, I. *et al.* (2010) Triptan-induced latent sensitization: a possible basis for medication overuse headache. *Ann Neurol* 67: 325–337.
- Di Lorenzo, C., Di Lorenzo, G., Sances, G., Ghiotto, N., Guaschino, E., Grieco, G. *et al.* (2009) Drug consumption in medication overuse headache is influenced by brain-derived neurotrophic factor

- Val66Met polymorphism. *J Headache Pain* 10: 349–355.
- Di Lorenzo, C., Sances, G., Di Lorenzo, G., Rengo, C., Ghiotto, N., Guaschino, E. *et al.* (2007) The Wolframín His611Arg polymorphism influences medication overuse headache. *Neurosci Lett* 424: 179–184.
- Diener, H. (2012) Detoxification for medication overuse headache is not necessary. *Cephalalgia* 32: 423–427.
- Diener, H., Bussone, G., Van Oene, J., Lahaye, M., Schwalen, S., Goadsby, P. *et al.* (2007) Topiramate reduces headache days in chronic migraine: a randomized, double-blind, placebo-controlled study. *Cephalalgia* 27: 814–823.
- Diener, H., Dichgans, J., Scholz, E., Geiselhart, S., Gerber, W. and Bille, A. (1989) Analgesic-induced chronic headache: long-term results of withdrawal therapy. *J Neurol* 236: 9–14.
- Diener, H., Dodick, D., Aurora, S., Turkel, C., Degryse, R., Lipton, R. *et al.* (2010) OnabotulinumtoxinA for treatment of chronic migraine: results from the double-blind, randomized, placebo-controlled phase of the Preempt 2 trial. *Cephalalgia* 30: 804–814.
- Diener, H., Dodick, D., Goadsby, P., Bigal, M., Bussone, G., Silberstein, S. *et al.* (2009) Utility of topiramate for the treatment of patients with chronic migraine in the presence or absence of acute medication overuse. *Cephalalgia* 29: 1021–1027.
- Diener, H. and Limmroth, V. (2004) Medication-overuse headache: a worldwide problem. *Lancet Neurol* 3: 475–483.
- Dodick, D. (2002) Debate: Analgesic overuse is a cause, not consequence, of chronic daily headache. Analgesic overuse is not a cause of chronic daily headache. *Headache* 42: 547–554.
- Dodick, D., Turkel, C., Degryse, R., Aurora, S., Silberstein, S., Lipton, R. *et al.* (2010) OnabotulinumtoxinA for treatment of chronic migraine: pooled results from the double-blind, randomized, placebo-controlled phases of the Preempt Clinical Program. *Headache* 50: 921–936.
- Dyb, G., Holmen, T. and Zwart, J. (2006) Analgesic overuse among adolescents with headache: The Head-Hunt-Youth Study. *Neurology* 66: 198–210.
- Eggen, A. (1993) The Tromso Study: frequency and predicting factors of analgesic drug use in a free-living population (12–56 years). *J Clin Epidemiol* 46: 1297–1304.
- Evers, S., Afra, J., Frese, A., Goadsby, P., Linde, M., May, A. *et al.* (2009) EFNS Guideline on the drug treatment of migraine – revised report of an EFNS task force. *Eur J Neurol* 16: 968–981.
- Evers, S. and Jensen, R. (2011) treatment of medication overuse headache – guideline of the EFNS Headache Panel. *Eur J Neurol* 18: 1115–1121.
- Evers, S. and Marziniak, M. (2010) Clinical features, pathophysiology, and treatment of medication-overuse headache. *Lancet Neurol* 9: 391–401.
- Ferrari, A., Cicero, A., Bertolini, A., Leone, S., Pasciullo, G. and Sternieri, E. (2006) Need for analgesics/drugs of abuse: a comparison between headache patients and addicts by the Leeds Dependence Questionnaire (LDQ). *Cephalalgia* 26: 187–193.
- Ferraro, S., Grazi, L., Muffatti, R., Nava, S., Ghielmetti, F., Bertolino, N. *et al.* (2012) In medication-overuse headache, FMRI shows long-lasting dysfunction in midbrain areas. *Headache* 52: 1520–1534.
- Fritsche, G., Frettlöh, J., Huppe, M., Dlugaj, M., Matatko, N., Gaul, C. *et al.* (2010) Prevention of medication overuse in patients with migraine. *Pain* 151: 404–413.
- Fuh, J. and Wang, S. (2012) Dependent behavior in patients with medication-overuse headache. *Curr Pain Headache Rep* 16: 73–79.
- Fuh, J., Wang, S., Lu, S. and Juang, K. (2005) Does medication overuse headache represent a behavior of dependence? *Pain* 119: 49–55.
- Fumal, A., Laureys, S., Di Clemente, L., Boly, M., Bohotin, V., Vandenheede, M. *et al.* (2006) Orbitofrontal cortex involvement in chronic analgesic-overuse headache evolving from episodic migraine. *Brain* 129: 543–550.
- Gaul, C., Bromstrup, J., Fritsche, G., Diener, H. and Katsarava, Z. (2011) Evaluating integrated headache care: a one-year follow-up observational study in patients treated at the Essen Headache Centre. *BMC Neurol* 11: 124.
- Ghiotto, N., Sances, G., Galli, F., Tassorelli, C., Guaschino, E., Sandrini, G. *et al.* (2009) Medication overuse headache and applicability of the ichd-ii diagnostic criteria: 1-year follow-up study (Care I Protocol). *Cephalalgia* 29: 233–243.
- Gossop, M., Darke, S., Griffiths, P., Hando, J., Powis, B., Hall, W. *et al.* (1995) The Severity of Dependence Scale (SDS): psychometric properties of the SDS in English and Australian samples of heroin, cocaine and amphetamine users. *Addiction* 90: 607–614.
- Grande, R., Aaseth, K., Benth, J., Gulbrandsen, P., Russell, M. and Lundqvist, C. (2009) The severity of dependence scale detects people with medication


- overuse: The Akershus Study of Chronic Headache. *J Neurol Neurosurg Psychiatry* 80: 784–789.
- Grande, R., Aaseth, K., Benth, J., Lundqvist, C. and Russell, M. (2011) Reduction in Medication-overuse headache after short information. The Akershus Study of Chronic Headache. *Eur J Neurol* 18: 129–137.
- Grande, R., Aaseth, K., Gulbrandsen, P., Lundqvist, C. and Russell, M. (2008) Prevalence of primary chronic headache in a population-based sample of 30- to 44-year-old persons. The Akershus Study of Chronic Headache. *Neuroepidemiology* 30: 76–83.
- Grazzi, L., Andrasik, F., D'amico, D., Usai, S., Kass, S. and Bussone, G. (2004) Disability in chronic migraine patients with medication overuse: treatment effects at 1-year follow-up. *Headache* 44: 678–683.
- Grazzi, L., Andrasik, F., Usai, S. and Bussone, G. (2008) In-patient versus day-hospital Withdrawal treatment for chronic migraine with medication overuse and disability assessment: results at one-year follow-up. *Neurol Sci* 29 Suppl 1: S161–S163.
- Grazzi, L. and Bussone, G. (2012) Medication overuse headache (MOH): complication of migraine or secondary headache? *Neurol Sci* 33 Suppl 1: S27–28.
- Hagen, K., Jensen, R., Boe, M. and Stovner, L. (2010) Medication overuse headache: a critical review of end points in recent follow-up studies. *J Headache Pain* 11: 373–377.
- Hagen, K., Linde, M., Steiner, T., Stovner, L. and Zwart, J. (2012) Risk factors for medication-overuse headache: an 11-year follow-up study. The Nord-Trøndelag Health Studies. *Pain* 153: 56–61.
- Hagen, K., Vatten, L., Stovner, L., Zwart, J., Krokstad, S. and Bovim, G. (2002) Low Socio-economic status is associated with increased risk of frequent headache: a prospective study of 22718 adults in Norway. *Cephalalgia* 22: 672–679.
- Hargreave, M., Andersen, T., Nielsen, A., Munk, C., Liaw, K. and Kjaer, S. (2010) Factors associated with a continuous regular analgesic use – a population-based study of more than 45,000 Danish women and men 18-45 years of age. *Pharmacoepidemiol Drug Saf* 19: 65–74.
- Headache Classification Committee of the International Headache Society (IHS) (2013) The International Classification of Headache Disorders, 3rd Edition (beta version). *Cephalalgia* 33: 629–808.
- Horton, B. and Peters, G. (1963) Clinical manifestations of excessive use of ergotamine preparations and magement of withdrawal effect: report of 52 cases. *Headache* 2: 214–227.
- Jensen, R. and Stovner, L. (2008) Epidemiology and comorbidity of headache. *Lancet Neurol* 7: 354–361.
- Jonsson, P., Hedenrud, T. and Linde, M. (2011) Epidemiology of medication overuse headache in the general Swedish population. *Cephalalgia* 31: 1015–1022.
- Jonsson, P., Linde, M., Hensing, G. and Hedenrud, T. (2012) Sociodemographic differences in medication use, health-care contacts and sickness absence among individuals with medication-overuse headache. *J Headache Pain* 13: 281–290.
- Katsarava, Z., Fritsche, G., Muessig, M., Diener, H. and Limmroth, V. (2001) Clinical features of withdrawal headache following overuse of triptans and other headache drugs. *Neurology* 57: 1694–1698.
- Katsarava, Z. and Jensen, R. (2007) Medication-overuse headache: where are we now? *Curr Opin Neurol* 20: 326–330.
- Katsarava, Z., Limmroth, V., Finke, M., Diener, H. and Fritsche, G. (2003) Rates and predictors for relapse in medication overuse headache: a 1-year prospective Study. *Neurology* 60: 1682–1683.
- Katsarava, Z., Muessig, M., Dzagnidze, A., Fritsche, G., Diener, H. and Limmroth, V. (2005) Medication overuse headache: rates and predictors for relapse in a 4-year prospective study. *Cephalalgia* 25: 12–15.
- Katsarava, Z. and Obermann, M. (2013) Medication-overuse headache. *Curr Opin Neurol* 26: 276–81.
- Kristoffersen, E., Grande, R., Aaseth, K., Lundqvist, C. and Russell, M. (2012a) Management of primary chronic headache in the general population: the Akershus Study of Chronic Headache. *J Headache Pain* 13: 113–120.
- Kristoffersen, E., Lundqvist, C., Aaseth, K., Grande, R. and Russell, M. (2013) Management of secondary chronic headache in the general population: the Akershus Study of Chronic Headache. *J Headache Pain* 14: 5.
- Kristoffersen, E., Straand, J., Benth, J., Russell, M. and Lundqvist, C. (2012b) Study Protocol: brief intervention for medication overuse headache – a double-blinded cluster randomised parallel controlled trial in primary care. *BMC Neurol* 12: 70.
- Krymchantowski, A. and Moreira, P. (2003) Out-patient detoxification in chronic migraine: comparison of strategies. *Cephalalgia* 23: 982–993.
- Kudrow, L. (1982) Paradoxical effects of frequent analgesic use. *Adv Neurol* 33: 335–341.
- Lake, A., III (2006) Medication overuse headache: biobehavioral issues and solutions. *Headache* 46 Suppl 3: S88–97.
- Lantéri-Minet, M., Auray, J., El Hasnaoui, A., Dartigues, J., Duru, G., Henry, P. et al. (2003) Prevalence and description of chronic daily headache

- in the general population in France. *Pain* 102: 143–149.
- Lantéri-Minet, M., Duru, G., Mudge, M. and Cottrell, S. (2011) Quality of life impairment, disability and economic burden associated with chronic daily headache, focusing on chronic migraine with or without medication overuse: a systematic review. *Cephalalgia* 31: 837–850.
- Latinovic, R., Gulliford, M. and Ridsdale, L. (2006) Headache and migraine in primary care: consultation, prescription, and referral rates in a large population. *J Neurol Neurosurg Psychiatry* 77: 385–387.
- Lauwerier, E., Paemeleire, K., Van Damme, S., Goubert, L. and Crombez, G. (2011) Medication use in patients with migraine and medication-overuse headache: the role of problem-solving and attitudes about pain medication. *Pain* 152: 1334–1339.
- Le Grand, S., Supornsilpchai, W., Saengjaroenatham, C. and Srikiatkachorn, A. (2011) Serotonin depletion leads to cortical hyperexcitability and trigeminal nociceptive facilitation *via* the nitric oxide pathway. *Headache* 51: 1152–1160.
- Limmroth, V., Katsarava, Z., Fritsche, G., Przywara, S. and Diener, H. (2002) Features of medication overuse headache following overuse of different acute headache drugs. *Neurology* 59: 1011–1014.
- Linde, M., Gustavsson, A., Stovner, L., Steiner, T., Barre, J., Katsarava, Z. *et al.* (2012) The Cost of headache disorders in Europe: The EuroLight Project. *Eur J Neurol* 19: 703–711.
- Linton-Dahlöf, P., Linde, M. and Dahlöf, C. (2000) Withdrawal therapy improves chronic daily headache associated with long-term misuse of headache medication: a retrospective study. *Cephalalgia* 20: 658–662.
- Lu, S., Fuh, J., Chen, W., Juang, K. and Wang, S. (2001) Chronic daily headache in Taipei, Taiwan: prevalence, follow-up and outcome predictors. *Cephalalgia* 21: 980–986.
- Lundqvist, C., Aaseth, K., Grande, R., Benth, J. and Russell, M. (2010) The severity of dependence score correlates with medication overuse in persons with secondary chronic headaches. The Akershus Study of Chronic Headache. *Pain* 148: 487–491.
- Lundqvist, C., Benth, J., Grande, R., Aaseth, K. and Russell, M. (2011) An adapted severity of dependence scale is valid for the detection of medication overuse: the Akershus Study of Chronic Headache. *Eur J Neurol* 18: 512–518.
- Lundqvist, C., Grande, R., Aaseth, K. and Russell, M. (2012) Dependence scores predict prognosis of medication overuse headache: a prospective cohort from the Akershus Study of Chronic Headache. *Pain* 153: 682–686.
- Mathew, N., Kurman, R. and Perez, F. (1990) Drug induced refractory headache—clinical features and management. *Headache* 30: 634–638.
- Mehuys, E., Paemeleire, K., Van Hees, T., Christiaens, T., Van Bortel, L., Van Tongelen, I. *et al.* (2012) Self-medication of regular headache: a community pharmacy-based survey. *Eur J Neurol* 19: 1093–1099.
- Meng, I., Dodick, D., Ossipov, M. and Porreca, F. (2011) Pathophysiology of medication overuse headache: insights and hypotheses from preclinical studies. *Cephalalgia* 31: 851–860.
- Meskunas, C., Tepper, S., Rapoport, A., Sheftell, F. and Bigal, M. (2006) Medications associated with probable medication overuse headache reported in a tertiary care headache center over a 15-year period. *Headache* 46: 766–772.
- Munksgaard, S., Bendtsen, L. and Jensen, R. (2012) Detoxification of medication-overuse headache by a multidisciplinary treatment programme is highly effective: a comparison of two consecutive treatment methods in an open-label design. *Cephalalgia* 32: 834–844.
- Munksgaard, S., Bendtsen, L. and Jensen, R. (2013) Modulation of central sensitisation by detoxification in MOH: results of a 12-month detoxification study. *Cephalalgia*:
- Olesen, J. (2012) Detoxification for medication overuse headache is the primary task. *Cephalalgia* 32: 420–422.
- Paemeleire, K., Bahra, A., Evers, S., Matharu, M. and Goadsby, P. (2006) Medication-overuse headache in patients with cluster headache. *Neurology* 67: 109–113.
- Patterson, V. and Esmonde, T. (1993) Comparison of the handling of neurological outpatient referrals by general physicians and a neurologist. *J Neurol Neurosurg Psychiatry* 56: 830–830.
- Perrotta, A., Serrao, M., Sandrini, G., Burstein, R., Sances, G., Rossi, P. *et al.* (2010) Sensitisation of spinal cord pain processing in medication overuse headache involves supraspinal pain control. *Cephalalgia* 30: 272–284.
- Peters, G. and Horton, B. (1950) Headache; with special reference to the excessive use of ergotamine tartrate and dihydroergotamine. *J Lab Clin Med* 36: 972–973.
- Peters, G. and Horton, B. (1951) Headache: with special reference to the excessive use of ergotamine preparations and withdrawal effects. *Proc Staff Meet Mayo Clin* 26: 153–161.
- Porteous, T., Bond, C., Hannaford, P. and Sinclair, H. (2005) How and why are non-prescription analgesics used in Scotland? *Fam Pract* 22: 78–85.

- Rabe, K., Pageler, L., Gaul, C., Lampl, C., Kraya, T., Foerderreuther, S. *et al.* (2013) Prednisone for the treatment of withdrawal headache in patients with medication overuse headache: a randomized, double-blind, placebo-controlled study. *Cephalalgia* 33: 202–207.
- Radat, F., Creac'h, C., Guegan-Massardier, E., Mick, G., Guy, N., Fabre, N. *et al.* (2008) Behavioral dependence in patients with medication overuse headache: a cross-sectional study in consulting patients using the DSM-IV criteria. *Headache* 48: 1026–1036.
- Radat, F., Creac'h, C., Swendsen, J., Lafittau, M., Irachabal, S., Dousset, V. *et al.* (2005) Psychiatric comorbidity in the evolution from migraine to medication Overuse Headache. *Cephalalgia* 25: 519–522.
- Radat, F. and Lanteri-Minet, M. (2010) What is the role of dependence-related behavior in medication-overuse headache? *Headache* 50: 1597–1611.
- Ridsdale, L., Clark, L., Dowson, A., Goldstein, L., Jenkins, L., Mccrone, P. *et al.* (2007) How do patients referred to neurologists for headache differ from those managed in primary care? *Br J Gen Pract* 57: 388–395.
- Rossi, C., Pini, L., Cupini, M., Calabresi, P. and Sarchielli, P. (2008a) Endocannabinoids in platelets of chronic migraine patients and medication-overuse headache patients: relation with serotonin levels. *Eur J Clin Pharmacol* 64: 1–8.
- Rossi, P., Di, L., Faroni, J., Cesarino, F., Nappi, G. and Di Lorenzo, C. (2006) Advice alone vs. structured detoxification programmes for medication overuse headache: a prospective, randomized, open-label trial in transformed migraine patients with low medical needs. *Cephalalgia* 26: 1097–1105.
- Rossi, P., Faroni, J. and Nappi, G. (2008b) Medication overuse headache: predictors and rates of relapse in migraine patients with low medical needs. A 1-year prospective study. *Cephalalgia* 28: 1196–1200.
- Rossi, P., Faroni, J. and Nappi, G. (2011) Short-term effectiveness of simple advice as a withdrawal strategy in simple and complicated medication overuse headache. *Eur J Neurol* 18: 396–401.
- Rossi, P., Faroni, J., Tassorelli, C. and Nappi, G. (2013) Advice alone *versus* structured detoxification programmes for complicated medication overuse headache (MOH): a prospective, randomized, open-label trial. *J Headache Pain* 14: 10.
- Rossi, P., Jensen, R., Nappi, G., Allena, M. and Consortium, C. (2009) A narrative review on the management of medication overuse headache: the steep road from experience to evidence. *J Headache Pain* 10: 407–417.
- Russell, M. and Lundqvist, C. (2012) Prevention and management of medication overuse headache. *Curr Opin Neurol* 25: 290–295.
- Sandrini, G., Perrotta, A., Tassorelli, C., Torelli, P., Brighina, F., Sances, G. *et al.* (2011) Botulinum toxin type-A in the prophylactic treatment of medication-overuse headache: a multicenter, double-blind, randomized, placebo-controlled, parallel group study. *J Headache Pain* 12: 427–433.
- Saper, J., Hamel, R. and Lake, A. III (2005) Medication overuse headache (MOH) is a biobehavioural disorder. *Cephalalgia* 25: 545–546.
- Saper, J. and Lake, A. III (2006) Medication overuse headache: type I and type II. *Cephalalgia* 26: 1262.
- Sarchielli, P., Rainero, I., Coppola, F., Rossi, C., Mancini, M., Pinessi, L. *et al.* (2008) Involvement of corticotrophin-releasing factor and orexin-A in chronic migraine and medication-overuse headache: findings from cerebrospinal fluid. *Cephalalgia* 28: 714–722.
- Scher, A., Lipton, R., Stewart, W. and Bigal, M. (2010) Patterns of medication use by chronic and episodic headache sufferers in the general population: results from the Frequent Headache Epidemiology Study. *Cephalalgia* 30: 321–328.
- Srikiatkachorn, A. and Anthony, M. (2006) Serotonin receptor adaptation in patients with analgesic-induced headache. *Cephalalgia* 16: 419–422.
- Srikiatkachorn, A., Maneesri, S., Govitrapong, P. and Kasantikul, V. (1998) Derangement of serotonin system in migrainous patients with analgesic abuse headache: clues from platelets. *Headache* 38: 43–49.
- Stovner, L., Hagen, K., Jensen, R., Katsarava, Z., Lipton, R., Scher, A. *et al.* (2007) The global burden of headache: a documentation of headache prevalence and disability worldwide. *Cephalalgia* 27: 193–210.
- Straube, A., Pfaffenrath, V., Ladwig, K., Meisinger, C., Hoffmann, W., Fendrich, K. *et al.* (2010) Prevalence of chronic migraine and medication overuse headache in Germany – the German DMKG Headache Study. *Cephalalgia* 30: 207–213.
- Sun-Edelstein, C., Bigal, M. and Rapoport, A. (2009) Chronic migraine and medication overuse headache: clarifying the Current International Headache Society classification criteria. *Cephalalgia* 29: 445–452.
- Supornsilpchai, W., Le Grand, S. and Srikiatkachorn, A. (2010a) Cortical hyperexcitability and mechanism of medication-overuse headache. *Cephalalgia* 30: 1101–1109.
- Supornsilpchai, W., Le Grand, S. and Srikiatkachorn, A. (2010b) Involvement of pro-nociceptive 5-HT_{2a} receptor in the pathogenesis of medication-overuse headache. *Headache* 50: 185–197.

- Supornsilpchai, W., Sanguanrangsirikul, S., Maneesri, S. and Srikiatkachorn, A. (2006) Serotonin depletion, cortical spreading depression, and trigeminal nociception. *Headache* 46: 34–39.
- Tepper, S. (2002) Debate: Analgesic overuse is a cause, not consequence, of chronic daily headache. Analgesic overuse is a cause of chronic daily headache. *Headache* 42: 543–547.
- Vos, T., Flaxman, A., Naghavi, M., Lozano, R., Michaud, C., Ezzati, M. *et al.* (2012) Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: A Systematic Analysis for the Global Burden of Disease Study 2010. *Lancet* 380: 2163–2196.
- Wang, S., Fuh, J., Lu, S. and Juang, K. (2006) Chronic daily headache in adolescents: prevalence, impact, and medication overuse. *Neurology* 66: 193–197.
- Wang, S., Fuh, J., Lu, S., Liu, C., Hsu, L., Wang, P. *et al.* (2000) Chronic daily headache in chinese elderly: prevalence, risk factors, and biannual follow-up. *Neurology* 54: 314–319.
- Wiendels, N., Knuistingh Neven, A., Rosendaal, F., Spinhoven, P., Zitman, F., Assendelft, W. *et al.* (2006) Chronic frequent headache in the general population: prevalence and associated factors. *Cephalalgia* 26: 1434–1442.
- Wilkinson, S., Becker, W. and Heine, J. (2001) Opiate use to control bowel motility may induce chronic daily headache in patients with migraine. *Headache* 41: 303–309.
- Zeeberg, P., Olesen, J. and Jensen, R. (2006a) Discontinuation of medication overuse in headache patients: recovery of therapeutic responsiveness. *Cephalalgia* 26: 1192–1198.
- Zeeberg, P., Olesen, J. and Jensen, R. (2006b) Probable medication-overuse headache: the effect of a 2-month drug-free period. *Neurology* 66: 1894–1898.
- Zidverc-Trajkovic, J., Pekmezovic, T., Jovanovic, Z., Pavlovic, A., Mijajlovic, M., Radojicic, A. *et al.* (2007) Medication overuse headache: clinical features predicting treatment outcome at 1-year follow-up. *Cephalalgia* 27: 1219–1225.
- Zwart, J., Dyb, G., Hagen, K., Ødegård, K., Dahl, A., Bovim, G. *et al.* (2003a) Depression and anxiety disorders associated with headache frequency. The Nord-Trøndelag Health Study. *Eur J Neurol* 10: 147–152.
- Zwart, J., Dyb, G., Hagen, K., Svebak, S., Holmen, J., Head, T. *et al.* (2003b) Analgesic use : a predictor of chronic pain and medication overuse headache. *Neurology* 61: 160–164.
- Zwart, J., Dyb, G., Hagen, K., Svebak, S., Stovner, L. and Holmen, J. (2004) Analgesic overuse among subjects with headache, neck, and low-back pain. *Neurology* 62: 1540–1544.

Visit SAGE journals online
<http://taw.sagepub.com>

 SAGE journals