

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/325203254>

The endocannabinoid–alcohol crosstalk: Recent advances on a bi–faceted target

Article in *Clinical and Experimental Pharmacology and Physiology* · May 2018

DOI: 10.1111/1440-1681.12967

CITATIONS

7

READS

166

6 authors, including:



Gianluca Lavanco

University of Catania

26 PUBLICATIONS 118 CITATIONS

[SEE PROFILE](#)



Valentina Castelli

Università degli Studi di Palermo

8 PUBLICATIONS 33 CITATIONS

[SEE PROFILE](#)



Anna Brancato

Università degli Studi di Palermo

54 PUBLICATIONS 678 CITATIONS

[SEE PROFILE](#)



Giuseppe Tringali

Catholic University of the Sacred Heart

98 PUBLICATIONS 2,444 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



Gambling [View project](#)



PROF. CARLA CANNIZZARO (Orcid ID : 0000-0002-7351-1712)

Article type : Review Article

Title:

The endocannabinoid-alcohol crosstalk: recent advances on a bi-faceted target

Authors:

Gianluca Lavanco¹, Valentina Castelli², Anna Brancato², Giuseppe Tringali³, Fulvio Plescia² and Carla Cannizzaro²

Affiliations

¹Department of Biomedical and Biotechnological Sciences, School of Medicine, University of Catania, Catania, Italy

²Department of Sciences for Health Promotion and Mother and Child Care "Giuseppe D'Alessandro", University of Palermo, Palermo, Italy

³Institute of Pharmacology, Catholic University School of Medicine, I 00168 Rome, Italy

Short Title

Endocannabinoid-alcohol crosstalk

Corresponding Author*

Carla Cannizzaro

Department of Sciences for Health Promotion and Mother and Child Care "G. D'Alessandro"

University of Palermo

Via del Vespro 129

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1111/1440-1681.12967

This article is protected by copyright. All rights reserved.

90127 Palermo, Italy

+39 0916555855

carla.cannizzaro@unipa.it

Abstract

Increasing evidence focuses on the endocannabinoid system as a relevant player in the induction of aberrant synaptic plasticity and related addictive phenotype following chronic excessive alcohol drinking. Besides, the endocannabinoid system is implicated in the pathogenesis of alcoholic liver disease. Interestingly, whereas the involvement of CB1 cannabinoid receptors in alcohol rewarding properties is established, the central and peripheral action of CB2 cannabinoid signalling is still to be elucidated. This review aims at giving the input to deepen knowledge on the role of the endocannabinoid system, highlighting the advancing evidence that suggests that CB1 and CB2 receptors may play opposite roles in the regulation of both the reinforcing properties of alcohol in the brain and the mechanisms responsible for cell injury and inflammation in the hepatic tissue. The manipulation of the endocannabinoid system could represent a bi-faceted strategy to counteract alcohol-related dysfunction in central transmission and liver structural and functional disarrangement.

Key words:

Alcohol; brain; CB1; CB2; endocannabinoids; liver

Introduction

Alcohol use disorder is a chronic relapsing disease characterized by compulsive alcohol seeking and taking¹, loss of control in limiting alcohol intake despite serious negative consequences, and recurring episodes of abstinence and relapse.²

According to the World Health Organization Report on Alcohol and Health (2011), alcohol abuse is responsible for at least 60 major types of systemic diseases,³ such as gut-derived inflammation⁴, increased risk of colorectal cancer^{5,6} cardiovascular and heart disease.^{7,8} However, the liver has been for long time considered the major victim of the harmful use of alcohol, since it is the main organ responsible for metabolizing alcohol. Several hypotheses have been advanced to explain the pathogenic mechanisms of alcohol-related liver disease, as well as of the aberrant functioning of the brain of alcoholics.⁹⁻¹¹ Accordingly, it would be overambitious, and out of our aim, to elucidate how neurotransmitters and modulators are affected by alcohol, contributing, in turn, to its consequences. Rather, this review will focus on the endocannabinoid system (ECS), as a common critical player in mediating both synaptic neuroadaptation and liver architectural distortion that result from excessive alcohol exposure. Indeed, the ECS is implicated in the regulation of a range of physiological processes and pathological conditions both at the central level and in periphery,¹²⁻¹⁴ including neural development, synaptic plasticity, pain, emotionality, immune function, metabolism and energy homeostasis.¹⁵⁻²² Moreover, if endocannabinoid(s) (eCB) involvement in the central effects of alcohol has been largely reported,^{23,24} some evidence has appeared about the role eCB play in mediating alcohol activity in the liver. This is a critical issue, because the identification of a common target accountable for alcohol-related brain and liver disease could pave the way for new therapeutic strategies.

Results

ECS-mediated Alcohol reinforcing effects

Alcohol produces its effects through actions on multiple brain circuits and involves neuroadaptive changes not only in adulthood but especially in critical periods of development.^{22,25-31}

The past two decades of clinical research, as well as data derived from preclinical models of alcohol addiction, point to the glutamatergic and the GABAergic systems as the main target of alcohol activity in the mesocorticolimbic system,³²⁻³⁸ where alcohol intake results in increased dopamine (DA) release from the ventral tegmental area (VTA) to the nucleus accumbens (NAc). Notwithstanding, alcohol's mechanism of action at the molecular level is fairly unknown. Some evidence of a direct effect on the dopaminergic system came from findings on the role played by acetaldehyde (ACD), its first oxidative metabolite. Indeed, ACD directly affects dopaminergic neurotransmission, increasing neuronal firing in the VTA, and stimulating DA release in the NAc shell.^{21,39-41} Moreover, ACD is able to induce and maintain an addictive-behavior in which seeking and relapse are modulated by the eCB^{12,42} so that ACD is believed to play a primary role in the "first-hit" of alcohol reinforcement and in the induction of relapse.^{21,22,39,43-46} Indeed, the ECS²¹ contributes to alcohol's rewarding effect, and to the occurrence of an addictive phenotype by fine tuning synaptic transmission: the eCB in fact, are key activity-dependent messengers that, by short and long-term decreases in synaptic transmission, regulate glutamatergic and GABAergic synapses. Long-term depression (LTD) is the best-characterized and widespread form of eCB mediated long-term synaptic plasticity. Because behavioural adaptations rely on changes in synaptic strength, the cannabinoid receptor type 1 (CB₁ receptor)-activated-LTD^{17,47} represents a fundamental mechanism for making long-term changes to neural circuits and behaviour. Notably, eCB induce synaptic plasticity in the VTA: CB₁ receptor-induced LTD of excitatory inputs to VTA GABA interneurons is basically mediated by 2-arachidonylglycerol (2-AG), which is formed postsynaptically via glutamate metabotropic receptor activation.^{48,49} Alcohol itself can induce an enhancement of CB₁ receptor-

dependent LTD by increasing eCB levels: as a matter of fact, alcohol is able to activate phospholipase A2 (PLA2), promote the synthesis of cannabinoid ligands arachidonylethanolamine (AEA) and 2-AG in neuronal cells, hinder fatty-acid amide hydrolase (FAAH) and monoacylglycerol lipase (MAGL) activity.^{50,51} This potentiation of CB₁ receptor-dependent LTD can consequently promote an enhancement in VTA dopaminergic signalling and result in increased alcohol rewarding effects. Interestingly, cue-induced alcohol craving can also result in a marked elevation of circulating levels of AEA,⁵² suggesting that AEA mobilization in peripheral tissues may be a marker of activation of the ECS in the brain.

The involvement of CB₁ receptors in alcohol rewarding effect is further confirmed by studies showing that the administration of CB₁ receptor agonists increases alcohol consumption in animal models of addiction.^{53,54} In particular, the administration of WIN 55,212-2, a synthetic non-selective CB₁ receptor agonist, causes a long-lasting increase in alcohol consumption during the re-exposure to alcohol in an animal model of relapse-like drinking⁵⁵ and is able to modulate binge-like alcohol intake in male mice.⁵⁶ Moreover, CB₁ receptor agonist CP-55,940 increases alcohol preference in a two-bottle choice paradigm⁵⁷ and alcohol-maintained responses in a progressive ratio paradigm.⁵⁸ On the other hand, the systemic administration of CB₁ receptor antagonists, such as SR141716A (rimonabant) and AM251, is able to decrease voluntary alcohol intake in naïve^{59,60} and in alcohol-preferring rats and mice.^{61,62} CB₁ receptor antagonists can also reduce alcohol-seeking behaviour in a rat model of relapse⁶³ and decrease binge alcohol and sucrose consumption in adolescent and adult mice.⁵⁴ Overall, the role of CB₁ receptors on alcohol reinforcing properties has been mainly unveiled by research on CB₁ receptor knockout mice, whose receptor deletion significantly reduces alcohol preference⁶⁴ and decreases its intake and conditioned place preference.^{65,66,67} Interestingly, Hungund et al. showed that mice lacking CB₁ receptors consume less alcohol than their wild-type counterparts and do not display alcohol-induced increase in DA release in the NAc.⁶⁵

In contrast, Linsenbardt and Bohem⁵⁶ showed a decrease in alcohol intake in male mice induced by high doses of WIN 55,212-2 in the first 30 minutes of Drinking In the Dark; accordingly, Cippitelli et al.⁶⁸ reported a decline in alcohol self-administration in Wistar rats, as a consequence of CB₁ receptors stimulation. However, it cannot be ruled out that the reduction in fluid consumption may be due to the motor inhibition component of the CB₁ receptor agonism. Moreover, inconsistent results came from studies employing cannabinoid receptor deletion: indeed, CB₁ receptor-deficient mice initially show a higher alcohol preference and alcohol intake than wild type animals, while, after the first week, CB₁ receptor deficient mice display a similar drinking pattern as wild type, suggesting that cannabinoid modulation of the rewarding stimuli plays a fundamental role in alcohol approach.⁶⁹

Besides the involvement of classical neurotransmitters, the eCB's modulation of the response to alcohol involves the recruitment of reward/stress-related neuropeptides - such as CRH and Neuropeptide Y (NPY) – which play a critical role in the development of addiction.^{70,71} Indeed, recent findings show that rat exposure to high concentration of ACD induces an increase in eCB transmission and this results in a downregulation of NPY expression and the occurrence of withdrawal symptoms following ACD suspension.^{21,22} Moreover, the study showed that CB₁ receptor blockade by the antagonist/inverse agonist AM281 is able to increase NPY expression, decrease ACD seeking-behaviour and boost homeostatic functional recovery.^{21,22} This is in accordance with AEA and 2-AG effect on CB₁ receptors on glutamate terminals, and the resulting decrease in glutamate release and downregulation in NPY mRNA levels^{72,73}. Overall CB₁ receptor signalling in the mesocorticolimbic system is a fundamental prerequisite for the expression of motivation to seek rewarding stimuli (Figure 1a).

CB₂ receptors: A Novel Target for Addiction

When cannabinoid receptors type 2 (CB₂ receptor) were first cloned, they were tagged as peripheral receptors.⁷⁴ However, recent studies suggest that CB₂ receptors are expressed in the nervous system – certainly in activated microglia and in some neuron subsets.⁷⁵

Indeed, CB₂ receptors have been recently involved in synaptic plasticity,^{76,77} and latest studies investigated the role of CB₂ receptors in models of alcohol, nicotine and cocaine addiction⁷⁸⁻⁸⁰ suggesting a role, not well defined yet, for CB₂ receptors in the modulation of drug reward-related behaviours.⁸¹

A recent study by Liu and colleagues provided a powerful new genetic tool for elucidating the functional role of CB₂ receptors in the CNS, by their selective deletion in dopaminergic neurons.⁸²

They demonstrated that CB₂ receptors put a “brake” on the classical locomotor activation that follows dopaminergic stimulation, since their conditional deletion in DAT positive midbrain dopaminergic neurons in DAT-Cnr2 cKO mice enhances psychomotor behavior. Traditionally CB₁ receptor agonism was associated with the tetrad effects in mice, producing the characteristic profile of suppression of locomotion, antinociception, hypothermia and catalepsy. This notion had been supported using data from radioligand binding and in-vivo behavioral assays that lacked sensitivity and cell-type specific deletion of the CCB₂ receptors. The dopaminergic neuron-specific deletion of the CB₂ receptors allowed providing evidence of the contribution of both CB₁ receptors and CB₂ receptors in the cannabinoid tetrad task. This is consistent with preclinical studies showing that selective CB₂ receptor agonists exert an attenuation of neuropathic pain pathways in rodent pain models^{83,84} and suggest that CB₂ receptors may contribute to the physical effect resulting from ECS direct (or alcohol-evoked) activation.

Both CB₁ receptor and CB₂ receptor pathways show great diversity and complexity with distinct pre- or post-synaptic distribution patterns where they may work cooperatively or in opposition to modulate the effects of cannabinoid and eCB in diverse brain areas. Thus, activation of postsynaptic CB₂ receptors in relevant brain areas^{85,86,87} supports the inhibition of VTA DA neuronal firing by CB₂ receptors. This may be associated with the resistance of the DAT-Cnr2 cKO to the induction of conditioned place preference caused by alcohol. Presynaptic CB₁ receptors indeed are not expressed in VTA DA neurons and the major endocannabinoid function in DA neuron in the VTA is mediated by postsynaptic CB₂ receptors that play an inhibitory role with respect to alcohol's rewarding effect.

However, the DAT-Cnr2 cKO mice showed an increased sensitivity to the rewarding effects of cocaine and this contributes to the existing evidence regarding the role of CB₂ receptors in the addicted properties of this drug. Notably, the activation of brain CB₂ receptors inhibits cocaine self-administration, and increases in locomotion, DA neuronal firing and extracellular DA release in the NAc in mice.⁸⁸ This inhibitory effect is reversed by pharmacological blockade of CB₂ receptors by AM630 or is absent in CB₂ receptor knockout mice, suggesting specificity in CB₂ receptor-mediated effect. This apparent discrepancy, together with the modulating effect of alcohol reinforcing properties, illustrates the complexity of CB₂ receptors involvement in addiction.

Interestingly, CB₂ receptors are expressed not only on VTA dopaminergic neurons,^{88,89} but also in the hippocampus and substantia nigra where they occupy a postsynaptic position⁹⁰. These localizations might be associated with psychomotor and cognitive specific functions in balance with the activation of presynaptic CB₁ receptors.

Given that dysfunction in DA signalling is a major abnormality in psychiatric and neurological disorders,⁹¹ these studies provide a basis for further analysis of the role of CB₂ receptors in the etiology of central disorders associated with DA dysregulation and for the development of drugs that selectively target eCB receptors (Figure 1a).

ECS-mediated effects in Alcoholic Liver Disease

The ECS and its receptors have emerged as major regulators of several pathophysiological mechanisms responsible for cell injury and inflammatory response that are associated with chronic liver disease progression.⁹²⁻⁹⁴ Indeed, under physiological conditions, both hepatocytes and non-parenchymal cells (i.e. Kupffer) are able to produce eCB, but the basal expression of CB₁ receptors and CB₂ receptors in the adult healthy liver is low or even absent.^{95,96} Notably, eCB synthesis and the hepatic expression of both cannabinoid receptors are upregulated during chronic liver damage.⁹²

Studies applying genetic or pharmacological inactivation of cannabinoid receptors showed that CB₁ receptors and CB₂ receptors exert opposite effects on fibrogenesis: CB₁ receptors have profibrogenic effects,⁹⁵ while CB₂ receptors can inhibit or reverse hepatic fibrogenesis and exert anti-inflammatory effects,⁹⁷ thus protecting from liver damage progression (Figure 1b).^{14,98} Notably, alcohol intake increases the hepatic expression of CB₁ receptors and, in activated stellate cells, it upregulates 2-AG, which, in turn, stimulates the deposition of fat in neighbouring hepatocytes by binding overexpressed CB₁ receptors.⁹⁹ On the other hand, CB₁ receptor knockout mice are resistant to the steatogenic and fibrogenic effects of alcohol, while CB₂ receptor knockout mice display increased collagen deposition, liver fat, and enhanced inflammatory scores.⁹⁵⁻¹⁰⁰

Accordingly, selective antagonists and agonists of respectively CB₁ receptors and CB₂ receptors, may attenuate the development of alcoholic liver disease.^{99,101,102} In particular, CB₁ receptor blockade by rimonabant and AM6545 induces partial regression of fibrosis and steatosis, restoring liver architecture.¹⁰³ Rimonabant administration to wild-type mice, as well as the genetic inactivation of CB₁ receptors are both associated with a significant reduction in fibrosis progression⁹⁵, reduced hepatic expression of the profibrogenic cytokines and decreased number of fibrogenic cells. Antifibrogenic properties of the CB₁ receptor-selective antagonist were ascribed to the antiproliferative and apoptotic properties of the compound in hepatic myofibroblasts.¹⁰⁴ Notwithstanding the beneficial effects of rimonabant in ameliorating hepatic fibrosis, the compound

was withdrawn from the market because of its psychotropic side effects, such as mood disorders that, in susceptible individuals, could also lead to major depression with high suicidal risk.¹⁰⁵

However novel compounds are currently under investigation in preclinical settings, able to modulate the eCB signalling while sparing the brain from toxicity.

Indeed, Denaës and colleagues recently demonstrated that CB₂ receptors agonism can protect from alcoholic liver disease by inhibiting hepatic inflammation and steatosis in Kupffer cell through an autophagy-dependent pathway.¹⁰² Moreover, CB₂ receptor agonism can facilitate hepatic regeneration, as well as normalization of serum liver enzymes, triglycerides, free fatty acids, and cholesterol.¹⁰³ In alcohol-exposed mice, endogenous or exogenous activation of CB₂ receptors prevents Kupffer cells from switching into a pro-inflammatory phenotype and hepatocytes from accumulating triglycerides.⁹⁷ In vitro activation of CB₂ receptors regulates macrophage polarization by preventing pro-inflammatory responses and inducing polarization towards an anti-inflammatory phenotype.⁹⁷ Because CB₂ receptors are not expressed in hepatocytes,¹⁰⁶ the antisteatogenic signal could originate from Kupffer cells.

Overall these data pinpoint on the CB₁ receptor signalling pathway as a novel paradigm to explain the molecular mechanisms underlying alcohol-induced liver damage. Moreover, hepatic CB₂ receptors emerge as protective targets not only for their antifibrogenic properties but also for promoting hepatocyte survival and regeneration.

The rising availability and use of legal cannabis by individuals who also consume alcohol, make the assessment of cannabis impact on alcohol-related pathologies mandatory. Daily cannabis smoking has been pointed as an additive factor in the evolution of ALD.⁵ However, the synergistic effects of cannabis and alcohol on liver disease progression remains unclear and further research in this field is highly needed.

Discussion

The rapidly advancing evidence on the neurobiology of the ECS suggests that CB₁ receptors and CB₂ receptors may work independently and/or cooperatively to regulate diverse physiological functions and pathogenetic processes. Based on the present findings, we support the hypothesis of CB₁ receptors and CB₂ receptors playing opposite roles in the regulation of the reinforcing properties of alcohol that, in turn, is related to stimulation of eCB transmission at central and peripheral levels. The ECS regulates both pro- and antifibrogenic responses in the liver with CB₁ receptors exerting profibrogenic effects and CB₂ receptors inhibiting hepatic fibrogenesis and exerting anti-inflammatory effects. Overall ECS manipulation could represent a common strategy to reduce the incentive reinforcing properties of alcohol and to recover liver architecture and functional disarrangement. Further elucidation on the eCB-alcohol-crosstalk, will help finding specific agents able to respond to complex therapeutic demand.

Methods

The literature search targeted evidence-based guidelines, evidence-based summaries, systematic reviews and recent experimental research on alcohol and endocannabinoid central, peripheral and behavioural effects.

Acknowledgments

This research was supported by a 2012-ATE-0167 grant from the University of Palermo, Italy.

Disclosures

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

1. George O, Hope BT. Cortical and amygdalar neuronal ensembles in alcohol seeking, drinking and withdrawal. *Neuropharmacology*. 2017;122:107-114.
2. Koob GF. Theoretical frameworks and mechanistic aspects of alcohol addiction: alcohol addiction as a reward deficit disorder. *Curr. Top. Behav. Neurosci*. 2013;13, 3–30.
3. Papineni JK, Pinnelli VBK, Davanum R. Thyroid Hormone Levels in Chronic Alcoholic Liver Disease Patients Before and After Treatment. *J Clin Diagn Res*. 2017;11(7):BC13-BC16.
4. Bishehsari F, Magno E, Swanson G, Desai V, Voigt RM, Forsyth CB, Keshavarzian A. Alcohol and gut-derived inflammation. *Alcohol Res. Curr. Rev*. 2017;38:163–171.
5. Cai S, Li Y, Ding Y, Chen K, Jin M. Alcohol drinking and the risk of colorectal cancer death: A meta-analysis. *Eur. J. Cancer Prev*. 2014;23:532–539.
6. Hagggar FA, Boushey RP. Colorectal cancer epidemiology: Incidence, mortality, survival, and risk factors. *Clin. Colon Rectal Surg*. 2009;22:191–197.
7. Moore RD, Pearson TA. Moderate alcohol consumption and coronary artery disease: a review. *Medicine (Baltimore)*. 1986;65:242-246.

8. Sano F, Ohira T, Kitamura A, et al. Heavy alcohol consumption and risk of atrial fibrillation. The Circulatory Risk in Communities Study (CIRCS). *Circ. J.* 2014;78(4):955–61.
9. Kim YJ, Hwang SH, Jia Y, Seo WD, Lee SJ. Barley sprout extracts reduce hepatic lipid accumulation in ethanol-fed mice by activating hepatic AMP-activated protein kinase. *Food Res. Int.* 2017;101:209-217.
10. Wang Y, Mukhopadhyay P, Cao Z, Wang H, Feng D, Haskó G, Mechoulam R, Gao B, Pacher P. Cannabidiol attenuates alcohol-induced liver steatosis, metabolic dysregulation, inflammation and neutrophil-mediated injury. *Sci. Rep.* 2017;7:12064.
11. Chan LN, Anderson GD. Pharmacokinetic and pharmacodynamic drug interactions with ethanol (alcohol). *Clin. Pharmacokinet.* 2014;53(12):1115-36.
12. Cavallaro, A, Martines, F, Cannizzaro, C, Lavanco, G, Brancato, A, Carollo, G, Plescia, F, Salvago, P, Cannizzaro, E, Mucia, M, Rizzo, S, Martini, A, Plescia, F. Role of cannabinoids in the treatment of tinnitus. *Acta Med. Mediterr.* 2016;32:463.
13. Bedse G, Romano A, Lavecchia AM, Cassano T, Gaetani S. The role of endocannabinoid signaling in the molecular mechanisms of neurodegeneration in Alzheimer's disease. *J Alzheimers Dis.* 2015;43(4):1115-36.
14. Tam J, Liu J, Mukhopadhyay B, Cinar R, Godlewski G, Kunos G. Endocannabinoids in liver disease. *Hepatology.* 2011;53(1):346-55.

15. Rizzo V, Ferraro G, Carletti F, Lonobile G, Cannizzaro C, Sardo P. Evidences of cannabinoids-induced modulation of paroxysmal events in an experimental model of partial epilepsy in the rat. *Neurosci. Lett.* 2009;462(2):135-9.
16. Viveros MP, de Fonseca FR, Bermudez-Silva FJ, McPartland JM. Critical role of the endocannabinoid system in the regulation of food intake and energy metabolism, with phylogenetic, developmental, and pathophysiological implications. *Endocr. Metab. Immune Disord. Drug Targets.* 2008;8(3):220-30.
17. Heifets BD, Castillo PE. Endocannabinoid signaling and long-term synaptic plasticity. *Annu. Rev. Physiol.* 2009;71:283-306.
18. Guindon J, Hohmann AG. The endocannabinoid system and pain. *CNS Neurol. Disord. Drug Targets.* 2009;8(6):403-21.
19. Lutz B. Endocannabinoid signals in the control of emotion. *Curr. Opin. Pharmacol.* 2009;9(1):46-52.
20. Brancato A, Lavanco G, Cavallaro A, Plescia F, Cannizzaro C. The use of the Emotional-Object Recognition as an assay to assess learning and memory associated to an aversive stimulus in rodents. *J. Neurosci. Methods.* 2016;274:106-115.
21. Plescia F, Brancato A, Marino RA, Cannizzaro C. Acetaldehyde as a drug of abuse: insight into AM281 administration on operant-conflict paradigm in rats. *Front. Behav. Neurosci.* 2013;7:64.

22. Plescia F, Brancato A, Marino RA, Vita C, Navarra M, Cannizzaro C. Effect of Acetaldehyde Intoxication and Withdrawal on NPY Expression: Focus on Endocannabinoidergic System Involvement. *Front. Psychiatry*. 2014;5:138.
23. Basavarajappa BS. Fetal Alcohol Spectrum Disorder: Potential Role of Endocannabinoids Signaling. *Brain Sci*. 2015;5(4):456-93.
24. Henderson-Redmond AN, Guindon J, Morgan DJ. Roles for the endocannabinoid system in ethanol-motivated behavior. *Prog. Neuropsychopharmacol. Biol. Psychiatry*. 2016;65:330-9. 33
25. Abrahao KP, Salinas AG, Lovinger DM. Alcohol and the Brain: Neuronal Molecular Targets, Synapses, and Circuits. *Neuron*. 2017;96(6):1223 – 1238.
26. Koob GF. Focus on: Neuroscience and treatment: the potential of neuroscience to inform treatment. *Alcohol Res. Health*. 2010;33(1-2):144-51.
27. Brancato A, Cannizzaro C. Mothering under the influence: how perinatal drugs of abuse alter the mother-infant interaction. *Rev. Neurosci*. 2017.
28. Brancato A, Plescia F, Lavanco G, Cavallaro A, Cannizzaro C. Continuous and Intermittent Alcohol Free-Choice from Pre-gestational Time to Lactation: Focus on Drinking Trajectories and Maternal Behavior. *Front. Behav. Neurosci*. 2016;10:31.
29. Sirohi S, Bakalkin G, Walker BM. Alcohol-induced plasticity in the dynorphin/kappa-opioid receptor system. *Front. Mol. Neurosci*. 2012;5:95.

30. Cacace S, Plescia F, Sardo P, Cannizzaro C. Alcohol preference, behavioural reactivity and cognitive functioning in female rats exposed to a three-bottle choice paradigm. *Behav. Brain Res.* 2012;234(1):11-9.
31. Cacace S, Plescia F, La Barbera M, Cannizzaro C. Evaluation of chronic alcohol self-administration by a 3-bottle choice paradigm in adult male rats. Effects on behavioural reactivity, spatial learning and reference memory. *Behav. Brain Res.* 2011;219(2):213-20.
32. Burnett EJ, Chandler LJ, Trantham-Davidson H. Glutamatergic plasticity and alcohol dependence-induced alterations in reward, affect and cognition. *Prog. Neuropsychopharmacol. Biol. Psychiatry.* 2016;65:309-20.
33. Bell RL, Hauser SR, McClintick J, Rahman S, Edenberg HJ, Szumlinski KK, McBride WJ. Ethanol-Associated Changes in Glutamate Reward Neurocircuitry: A Minireview of Clinical and Preclinical Genetic Findings. *Prog. Mol. Biol. Transl. Sci.* 2016;137:41-85.
34. Naassila M, Pierrefiche O. GluN2B Subunit of the NMDA Receptor: The Keystone of the Effects of Alcohol During Neurodevelopment. *Neurochem. Res.* 2018.
35. Castelli V, Brancato A, Cavallaro A, Lavanco G, Cannizzaro C. Homer2 and Alcohol: A Mutual Interaction. *Front. Psychiatry.* 2017;8:268.
36. Chandler CM, Overton JS, Rüedi-Bettschen D, Platt DM. GABA(A) Receptor Subtype Mechanisms and the Abuse-Related Effects of Ethanol: Genetic and Pharmacological Evidence. *Handb. Exp. Pharmacol.* 2017.

37. Lindemeyer AK, Shen Y, Yazdani F, Shao XM, Spigelman I, Davies DL, Olsen RW, Liang J. α 2 Subunit-Containing GABA(A) Receptor Subtypes Are Upregulated and Contribute to Alcohol-Induced Functional Plasticity in the Rat Hippocampus. *Mol. Pharmacol.* 2017;92(2):101-112.
38. Weiner JL, Valenzuela CF. Ethanol modulation of GABAergic transmission: the view from the slice. *Pharmacol. Ther.* 2006;111(3):533-54.
39. Brancato A, Plescia F, Marino RA, Maniaci G, Navarra M, Cannizzaro C. Involvement of dopamine D2 receptors in addictive-like behaviour for acetaldehyde. *PLoS One.* 2014;13;9(6):e99454.
40. Sutura FM, De Caro V, Cannizzaro C, Giannola LI, Lavanco G, Plescia F. Effects of DA-Phen, a dopamine-aminoacidic conjugate, on alcohol intake and forced abstinence. *Behav. Brain Res.* 2016;310:109-18.
41. Spiga S, Talani G, Mulas G, Licheri V, Fois GR, Muggironi G, Masala N, Cannizzaro C, Biggio G, Sanna E, Diana M. Hampered long-term depression and thin spine loss in the nucleus accumbens of ethanol-dependent rats. *Proc. Natl. Acad. Sci. U S A.* 2014;111(35):E3745-54.
42. Brancato A, Lavanco G, Cavallaro A, Plescia F, Cannizzaro C. Acetaldehyde, Motivation and Stress: Behavioral Evidence of an Addictive ménage à trois. *Front. Behav. Neurosci.* 2017;9:11:23.
43. Israel Y, Quintanilla ME, Karahanian E, Rivera-Meza M, Herrera-Marschitz M. The "first hit" toward alcohol reinforcement: role of ethanol metabolites. *Alcohol Clin. Exp. Res.* 2015;39(5):776-86.

44. Plescia F, Brancato A, Venniro M, Maniaci G, Cannizzaro E, Sutera FM, De Caro V, Giannola LI, Cannizzaro C. Acetaldehyde self-administration by a two-bottle choice paradigm: consequences on emotional reactivity, spatial learning, and memory. *Alcohol*. 2015;49(2):139-48.
45. Cannizzaro C, La Barbera M, Plescia F, Cacace S, Tringali G. Ethanol modulates corticotropin releasing hormone release from the rat hypothalamus: does acetaldehyde play a role? *Alcohol Clin. Exp. Res*. 2010;34(4):588-93. doi: 10.1111/j.1530-0277.2009.01127.x.
46. Cacace S, Plescia F, Barberi I, Cannizzaro C. Acetaldehyde oral self-administration: evidence from the operant-conflict paradigm. *Alcohol Clin. Exp. Res*. 2012;36(7):1278-87. doi: 10.1111/j.1530-0277.2011.01725.x.
47. Castillo PE, Younts TJ, Chávez AE, Hashimoto Y. Endocannabinoid signaling and synaptic function. *Neuron*. 2012;76(1):70-81.
48. Friend L, Weed J, Sandoval P, Nufer T, Ostlund I, Edwards JG. CB1-Dependent Long-Term Depression in Ventral Tegmental Area GABA Neurons: A Novel Target for Marijuana. *J. Neurosci*. 2017;37(45):10943-10954.
49. Lupica CR, Riegel AC. Endocannabinoid release from midbrain dopamine neurons: a potential substrate for cannabinoid receptor antagonist treatment of addiction. *Neuropharmacology*. 2005;48(8):1105-16.
50. Basavarajappa BS, Saito M, Cooper TB, Hungund BL. Stimulation of cannabinoid receptor agonist 2-arachidonylglycerol by chronic ethanol and its modulation by specific neuromodulators in cerebellar granule neurons. *Biochim. Biophys. Acta*. 2000;1535(1):78-86.

51. Basavarajappa BS, Saito M, Cooper TB, Hungund BL. Chronic ethanol inhibits the anandamide transport and increases extracellular anandamide levels in cerebellar granule neurons. *Eur. J. Pharmacol.* 2003;466(1-2):73-83.
52. Mangieri RA, Hong KI, Piomelli D, Sinha R. An endocannabinoid signal associated with desire for alcohol is suppressed in recently abstinent alcoholics. *Psychopharmacology (Berl)*. 2009;205(1):63-72.
53. Hungund BL, Basavarajappa BS. Role of endocannabinoids and cannabinoid CB1 receptors in alcohol-related behaviors. *Ann. N. Y. Acad. Sci.* 2004;1025:515-27.
54. Agolia AE, Holstein SE, Eastman VR, Hodge CW. Cannabinoid CB1 receptor inhibition blunts adolescent-typical increased binge alcohol and sucrose consumption in male C57BL/6J mice. *Pharmacol. Biochem. Behav.* 2016;143:11-7.
55. Alén F, Santos A, Moreno-Sanz G, González-Cuevas G, Giné E, Franco-Ruiz L, Navarro M, López-Moreno JA. Cannabinoid-induced increase in relapse-like drinking is prevented by the blockade of the glycine-binding site of N-methyl-D-aspartate receptors. *Neuroscience*. 2009;158(2):465-73.
56. Linsenbardt DN, Boehm SL 2nd. Agonism of the endocannabinoid system modulates binge-like alcohol intake in male C57BL/6J mice: involvement of the posterior ventral tegmental area. *Neuroscience*. 2009;164(2):424-34.
57. Colombo G, Serra S, Brunetti G, Gomez R, Melis S, Vacca G, Carai MM, Gessa L. Stimulation of voluntary ethanol intake by cannabinoid receptor agonists in ethanol-preferring sP rats. *Psychopharmacology (Berl)*. 2002;159(2):181-7.

58. Gallate JE, Saharov T, Mallet PE, McGregor IS. Increased motivation for beer in rats following administration of a cannabinoid CB1 receptor agonist. *Eur. J. Pharmacol.* 1999;370(3):233-40.
59. Femenía T, García-Gutiérrez MS, Manzanares J. CB1 receptor blockade decreases ethanol intake and associated neurochemical changes in fawn-hooded rats. *Alcohol Clin. Exp. Res.* 2010;34(1):131-41.
60. Freedland CS, Sharpe AL, Samson HH, Porrino LJ. Effects of SR141716A on ethanol and sucrose self-administration. *Alcohol Clin. Exp. Res.* 2001;25(2):277-82.
61. Colombo G, Agabio R, Fà M, Guano L, Lobina C, Loche A, Reali R, Gessa GL. Reduction of voluntary ethanol intake in ethanol-preferring sP rats by the cannabinoid antagonist SR-141716. *Alcohol Alcohol.* 1998;33(2):126-30.
62. Hungund BL, Basavarajappa BS, Vadasz C, Kunos G, Rodriguez de Fonseca F, Colombo G, Serra S, Parsons L, Koob GF. Ethanol, endocannabinoids, and the cannabinoidergic signaling system. *Alcohol Clin. Exp. Res.* 2002;26(4):565-74.
63. Cippitelli A, Bilbao A, Hansson AC, del Arco I, Sommer W, Heilig M, Massi M, Bermúdez-Silva FJ, Navarro M, Ciccocioppo R, de Fonseca FR; European TARGALC Consortium. Cannabinoid CB1 receptor antagonism reduces conditioned reinstatement of ethanol-seeking behavior in rats. *Eur. J. Neurosci.* 2005;21(8):2243-51.
64. Vinod KY, Kassir SA, Hungund BL, Cooper TB, Mann JJ, Arango V. Selective alterations of the CB1 receptors and the fatty acid amide hydrolase in the ventral striatum of alcoholics and suicides. *J. Psychiatr. Res.* 2010;44(9):591-7.

65. Hungund BL, Szakall I, Adam A, Basavarajappa BS, Vadasz C. Cannabinoid CB1 receptor knockout mice exhibit markedly reduced voluntary alcohol consumption and lack alcohol-induced dopamine release in the nucleus accumbens. *J. Neurochem.* 2003;84(4):698-704.

66. Naassila M, Pierrefiche O, Ledent C, Daoust M. Decreased alcohol self-administration and increased alcohol sensitivity and withdrawal in CB1 receptor knockout mice. *Neuropharmacology.* 2004;46(2):243-53.

67. Wang L, Liu J, Harvey-White J, Zimmer A, Kunos G. Endocannabinoid signaling via cannabinoid receptor 1 is involved in ethanol preference and its age-dependent decline in mice. *Proc. Natl. Acad. Sci. U S A.* 2003;100(3):1393-8.

68. Cippitelli A, Bilbao A, Gorriti MA, Navarro M, Massi M, Piomelli D, Ciccocioppo R, Rodríguez de Fonseca F. The anandamide transport inhibitor AM404 reduces ethanol self-administration. *Eur. J. Neurosci.* 2007;26(2):476-86

69. Ildiko Racz, Andras Bilkei-Gorzo, Zsuzsanna E. Toth, Kerstin Michel, Miklós Palkovits, Andreas Zimmer. A Critical Role for the Cannabinoid CB₁ Receptors in Alcohol Dependence and Stress-Stimulated Ethanol Drinking. *Journal of Neuroscience.* 2003;23(6):2453-2458.

70. Vengeliene V, Bilbao A, Molander A, Spanagel R. Neuropharmacology of alcohol addiction. *Br. J. Pharmacol.* 2008;154(2):299-315.

71. Olling JD, Ulrichsen J, Correll M, Woldbye DP. Gene expression in the neuropeptide Y system during ethanol withdrawal kindling in rats. *Alcohol Clin. Exp. Res.* 2010;34(3):462–7010.1111/j.1530-0277.2009.01110.x.

72. Cottone E, Guastalla A, Pomatto V, Campantico E, Palermo F, Magni AM, et al. Interplay of the endocannabinoid system with neuropeptide Y and corticotropin-releasing factor in the goldfish forebrain. *Ann. N. Y. Acad. Sci.* 2009;1163:372–510.
73. Misra K, Pandey SC. Differences in basal levels of CREB and NPY in nucleus accumbens regions between C57BL/6 and DBA/2 mice differing in inborn alcohol drinking behavior. *J. Neurosci. Res.* 2003;74(6):967–7510.
74. Munro S, Thomas KL, Abu-Shaar M. Molecular characterization of a peripheral receptor for cannabinoids. *Nature.* 1993;365(6441):61–65.
75. Atwood BK, Mackie K. CB2: A cannabinoid receptor with an identity crisis. *Br. J. Pharmacol.* 2010;160(3):467–479.
76. Stempel, AV et al. Cannabinoid type 2 receptors mediate a cell type-specific plasticity in the hippocampus. *Neuron.* 2016;90,795–809.
77. Onaivi, ES et al. CNS effects of CB2 cannabinoid receptors: beyond neuro-immuno-cannabinoid activity. *J. Psychopharmacol.* 2012;26:92–103.
78. Xi ZX, Peng XQ, Li X, Song R, Zhang HY, Liu QR, Yang HJ, Bi GH, Li J, Gardner EL. Brain cannabinoid CB₂ receptors modulate cocaine's actions in mice. *Nat. Neurosci.* 2011;14(9):1160-6.
79. Al Mansouri S, Ojha S, Al Maamari E, Al Ameri M, Nurulain SM, Bahi A. The cannabinoid receptor 2 agonist, β -caryophyllene, reduced voluntary alcohol intake and attenuated ethanol-induced place preference and sensitivity in mice. *Pharmacol. Biochem. Behav.* 2014;124:260-8.

80. Ignatowska-Jankowska BM, Muldoon PP, Lichtman AH, Damaj MI. The cannabinoid CB2 receptor is necessary for nicotine-conditioned place preference, but not other behavioral effects of nicotine in mice. *Psychopharmacology (Berl)*. 2013;229(4):591-601.
81. Ortega-Álvaro A, Ternianov A, Aracil-Fernández A, Navarrete F, García-Gutiérrez MS, Manzanares J. Role of cannabinoid CB2 receptor in the reinforcing actions of ethanol. *Addict. Biol*. 2015;20(1):43-55.
82. Liu QR, Canseco-Alba A, Zhang HY, Tagliaferro P, Chung M, Dennis E, Sanabria B, Schanz N, Escosteguy-Neto JC, Ishiguro H, Lin Z, Sgro S, Leonard CM, Santos-Junior JG, Gardner EL, Egan JM, Lee JW, Xi ZX, Onaivi ES. Cannabinoid type 2 receptors in dopamine neurons inhibits psychomotor behaviors, alters anxiety, depression and alcohol preference. *Sci. Rep*. 2017;7(1):17410.
83. Svizenska I.H. et al. Bilateral changes of cannabinoid receptor type 2 protein and mRNA in the dorsal root ganglia of a rat neuropathic pain model. *J. Histochem. Cytochem*. 2013;61:529–547.
84. Xu, J. et al. Activation of cannabinoid receptor 2 attenuates mechanical allodynia and neuroinflammatory responses in a chronic post-ischemic pain model of complex regional pain syndrome type 1 in rats. *Euro. J. Neurosc*. 2016;44:3046–3055.
85. Stempel AV, Stumpf A, Zhang HY, Özdoğan T, Pannasch U, Theis AK, Otte DM, Wojtalla A, Rácz I, Ponomarenko A, Xi ZX, Zimmer A, Schmitz D. Cannabinoid type 2 receptors mediate a cell type-specific plasticity in the hippocampus. *Neuron*. 2016;90:795–809.
86. Brusco A, Tagliaferro P, Saez T, Onaivi ES. Postsynaptic localization of CB2 cannabinoid receptors in the rat hippocampus. *Synapse*. 2008;62:944–949.

87. Brusco A, Tagliaferro PA, Saez T, Onaivi ES. Ultrastructural localization of neuronal brain CB2 cannabinoid receptors. *Ann. N. Y. Acad. Sci.* 2008;1139:450–457.

88. Zhang HY, Gao M, Liu QR, Bi GH, Li X, Yang HJ, Gardner EL, Wu J, Xi ZX. Cannabinoid CB2 receptors modulate midbrain dopamine neuronal activity and dopamine-related behavior in mice. *Proc- Natl. Acad. Sci. U S A.* 2014;111(46):E5007-15.

89. Zhang HY. et al. Expression of functional cannabinoid CB2 receptor in VTA dopamine neurons in rats. *Addiction Biology.* 2016.

90. Lanciego JL, Barroso-Chinea P, Rico AJ, Conte-Perales L, Callén L, Roda E, Gómez-Bautista V, López IP, Lluís C, Labandeira-García JL, Franco R. Expression of the mRNA coding the cannabinoid receptor 2 in the pallidal complex of *Macaca fascicularis*. *J Psychopharmacol.* 2011 Jan;25(1):97-104.

91. Maia TV and Frank MJ. From reinforcement learning models of the basal ganglia to the pathophysiology of psychiatric and neurological disorders. *Nat. Neurosci.* 2011;14:154–162.

92. Mallat A, Teixeira-Clerc F, Lotersztajn S. Cannabinoid signaling and liver therapeutics. *J. Hepatol.* 2013;59(4):891-6.

93. Patsenker E, Sachse P, Chicca A, Gachet MS, Schneider V, Mattsson J, Lanz C, Worni M, de Gottardi A, Semmo M, Hampe J, Schafmayer C, Brenneisen R, Gertsch J, Stickel F, Semmo N. Elevated levels of endocannabinoids in chronic hepatitis C may modulate cellular immune response and hepatic stellate cell activation. *Int. J. Mol. Sci.* 2015;16(4):7057-76.

94. Cao Z, Mulvihill MM, Mukhopadhyay P, et al. Monoacylglycerol lipase controls endocannabinoid and eicosanoid signaling and hepatic injury in mice. *Gastroenterology* 2013;144:808–17.
95. Teixeira-Clerc F, Julien B, Grenard P, Tran Van Nhieu J, Deveaux V, Li L, Serriere-Lanneau V, Ledent C, Mallat A, Lotersztajn S. CB1 cannabinoid receptor antagonists: a novel approach for the treatment of liver fibrosis. *Med. Sci. (Paris)*. 2006;22(8-9):683-5.
96. Julien B, Grenard P, Teixeira-Clerc F, Van Nhieu JT, Li L, Karsak M, Zimmer A, Mallat A, Lotersztajn S. Antifibrogenic role of the cannabinoid receptor CB2 in the liver. *Gastroenterology*. 2005;128(3):742-55.
97. Louvet A, Teixeira-Clerc F, Chobert MN, Deveaux V, Pavoine C, Zimmer A, Pecker F, Mallat A, Lotersztajn S. Cannabinoid CB2 receptors protect against alcoholic liver disease by regulating Kupffer cell polarization in mice. *Hepatology*. 2011;54(4):1217-26.
98. Pisanti S, Picardi P, Pallottini V, Martini C, Petrosino S, Proto MC, Vitale M, Laezza C, Gazzo P, Di Marzo V, Bifulco M. Anandamide drives cell cycle progression through CB1 receptors in a rat model of synchronized liver regeneration. *J. Cell. Physiol*. 2015;230(12):2905-14.
99. Jeong WI, Osei-Hyiaman D, Park O, Liu J, Bátkai S, Mukhopadhyay P, Horiguchi N, Harvey-White J, Marsicano G, Lutz B, Gao B, Kunos G. Paracrine activation of hepatic CB1 receptors by stellate cell-derived endocannabinoids mediates alcoholic fatty liver. *Cell. Metab*. 2008;7(3):227-35.
100. Trebicka J, Racz I, Siegmund SV, Cara E, Granzow M, Schierwagen R, Klein S, Wojtalla A, Hennenberg M, Huss S, Fischer HP, Heller J, Zimmer A, Sauerbruch T. Role of cannabinoid receptors in alcoholic hepatic injury: steatosis and fibrogenesis are increased in CB2 receptor-deficient mice and decreased in CB1 receptor knockouts. *Liver Int*. 2011;31(6):860-70.

101. Baldassarre M, Giannone FA, Napoli L, Tovoli A, Ricci CS, Tufoni M, Caraceni P. The endocannabinoid system in advanced liver cirrhosis: pathophysiological implication and future perspectives. *Liver Int.* 2013;33(9):1298-308.
102. Denaës T, Lodder J, Chobert MN, Ruiz I, Pawlotsky JM, Lotersztajn S, Teixeira-Clerc F. The Cannabinoid Receptor 2 Protects Against Alcoholic Liver Disease Via a Macrophage Autophagy-Dependent Pathway. *Sci. Rep.* 2016;6:28806.
103. Boesten JE, Kaper J, Stoffers HE, Kroon AA, van Schayck OC. Rimonabant improves obesity but not the overall cardiovascular risk and quality of life; results from CARDIO-REDUCE (CARDiometabolic Risk reDUCTiOn by Rimonabant: The Effectiveness in Daily practice and its USE). *Fam. Pract.* 2012;29(5):521-7.
104. Mallat A, Teixeira-Clerc F, Deveaux V, Manin S, Lotersztajn S. The endocannabinoid system as a key mediator during liver diseases: new insights and therapeutic openings. *Br. J. Pharmacol.* 2011;163(7):1432-40.
105. Moreira FA, Crippa JA. The psychiatric side-effects of rimonabant. *Rev. Bras. Psiquiatr.* 2009;31:145-53.
106. Teixeira-Clerc F, Belot MP, Manin S, Deveaux V, Cadoudal T, Chobert MN, Louvet A, Zimmer A, Tordjmann T, Mallat A, Lotersztajn S. Beneficial paracrine effects of cannabinoid receptor 2 on liver injury and regeneration. *Hepatology.* 2010;52(3):1046-59.

